DECLARATION

I, Dharitri Das, bearing Ph. D. Registration Number, Ph. D./1189/2010 dt

15.09.2010, hereby declare that the thesis entitled "Structure and reactivity of

ruthenium based anticancer drugs and their interaction with biomolecules: A

density functional approach" submitted to Assam University, Silchar for the partial

fulfillment of the degree of Doctor of Philosophy in Chemistry is the record of

original research work done by me under the supervision of Dr. Paritosh Mondal,

Department of Chemistry, Assam University, Silchar. I further declare that the

contents of this thesis have not formed the basis for the award of any other

degree/diploma.

Place: Assam University, Silchar

(Dharitri Das)

Date: 15.06.2015

Dedicated to My Father Tarun Chandra Das & Mother Ila Prava Das

ACKNOWLEDGEMENT

In the beginning, I owe it all to Almighty God for granting me the wisdom, health and strength to complete this research work.

I express my sincere gratitude to my supervisor, Dr. Paritosh Mondal for his invaluable guidance and constant encouragement during the thesis work. I am deeply indebted to him for the knowledge and experience I gained in theoretical field through his earnest efforts. I am very grateful for his active interest and fruitful discussions on several aspects of the thesis work. I thank him for his unfailing help and support throughout my Ph. D. programme.

I thank Prof. S. B. Paul, Prof. C. R. Bhattacharjee, Dr. P. C. Paul, Dr. Sk. Jasimuddin, Dr. M. K. Paul, Dr. D. Sengupta, Dr. H. Acharya, Dr. S. K. Ghosh, Dr. S. Choudhury, Dr. T. S. Singh and Dr. R. Panchadhayee, faculty members of the department for their support and encouragement throughout.

I extend my thanks to Mr. P. R Ramesh, Mr. S. Bhattacharjee, Mr. B. Nath, Mr. R. Kurmi as well as official staff members Md. J. Borbhuiya, Mr. L. Hmar and Mr. S. Bagdu for their cooperation.

I thank Mr. Samiran Garain, Ph. D. scholar, Jadavpur University, for providing journals which are not accessible from Assam University, whenever required.

I don't have enough words to express my thanks to my friend kalpana Upadhyaya and my brother Alok Ahmed Barbhuiya for their selfless help and support behind the screen.

I am thankful to my lab mates Abhijit Dutta, Dilip Chorei and all the research scholars of the department specially Sudip Pal and Zeaul Hoque Mazumder for their timely help and cooperation.

Words will be insufficient to thank my father Tarun Chandra Das and my mother Ila Prava Das for their encouragement at every stage of my life. My heartiest thank to them for being with me always. My greatest gratitude goes to my younger sister Bandita Das and my younger brother Bidyut Das who always support me in pursuing my studies. My family has always been there to support me through good and bad times and without their efforts I would not be able to reach at this stage in my life. I thank you all from the bottom of my heart.

I also like to acknowledge University Grants Commission (U.G.C), for financial support in the form of research fellowship.

I am thankful to Assam University, Silchar and the Department of Chemistry in particular for providing me the platform to complete my Ph. D programme.

Ph.D. is not the job of a single person. It is the work of all who supported you and tolerated you throughout this difficult period. I once again would like to thank all of you for being with me always.

(Dharitri Das)

ABBREVIATIONS

• FDA : American Food and Drugs Administration

• GG : guanine-guanine

• DNA : deoxyribonucleic acid

• DFT : density functional theory

• BLYP : Becke-Lee-Yang-Parr

• DNP : double numerical with polarization

• GGA : generalized gradient approximation

• LST : linear synchronous transit

• CG : conjugated gradient

• QST : quadratic synchronous transit

• IRC : intrinsic reaction coordinate

• MM : molecular mechanics

• QM : quantum mechanics

• QM/MM : quantum mechanics/molecular mechanics

• ONIOM : our Own N-layer Integrated molecular Orbital

molecular Mechanics

• GA : genetic algorithm

• apo-Tf : apo-transferrin

• QSAR : quantitative structure-activity relationship

• HT-29 : human Colorectal Adenocarcinoma Cell Line

• DMSO : dimethylsulfoxide

• im : imidazole

• IC_{50} : concentration of a compound that induces 50% growth

inhibition of cells compared to untreated cells

• CPCM : conductor-like polarizable continuum model

• his : histidine

• ser : serine

• cys : cysteine

• HSA : human serum albumin

• RCSB : research collaboratory for structural bioinformatics

• pdb : protein data bank

HOMO : highest occupied molecular orbitalLUMO : lowest unoccupied molecular orbital

• UFF : universal force field

• NO : nitric oxide

• ZPE : zero-point vibrational effects

• NBO : natural bond orbital

LIST OF TABLES Selected bond lengths (Å) and bond angles (⁰) calculated for Table 3.1 49 all complexes at B3LYP level both in the gas phase and solvent phase. Solvent phase calculated values are given in parenthesis Table 3.2 Calculated hardness (η in eV), chemical potential (μ in eV), 51 electrophilicity (ω in eV), local philicity (ω ⁺ in eV) for all complexes in gas phase and solvent phase. Local philicity value for the ruthenium atom is reported Selected bond lengths (Å) and bond angles (⁰) calculated for Table 4.1 61 all stationary points in the first step of Cl hydrolysis for the complexes II and IV Selected bond lengths (Å) and bond angles (⁰) calculated for Table 4.2 64 all stationary points in the second step of Cl hydrolysis of complex II and IV Selected bond lengths (Å) and bond angles (⁰) calculated for Table 4.3 67 all stationary points in the first step of DMSO hydrolysis of complex II and IV Selected bond lengths (Å) and bond angles (⁰) calculated for Table 4.4 68 all stationary points in the second step of DMSO hydrolysis of complex II and IV 70 Table 4.5 Total ZPE ($E_{tot/ZPE}$), Change of enthalpies (ΔH), Gibbs free energies (ΔG) and entropies (ΔS) at 298.15 K for all intermediate species and transition States of the hydrolysis processes of complex II Total ZPE ($E_{\it tot/ZPE}\,$), Change of enthalpies ($\Delta H\,$), Gibbs free 71 Table 4.6 energies (ΔG) and entropies (ΔS) at 298.15 K for all intermediate species and transition States of the hydrolysis processes of complex IV Selected bond lengths (Å) and bond angles (⁰) calculated for Table 5A.1 85 ruthenium(III) complexes at B3LYP level in the gas phase

- Table 5A.2 Energies of HOMO (E_H in eV) and LUMO (E_L in eV) and chemical hardness (η in eV) of ruthenium(III) complexes

 Table 5A.3 Binding energy (ΔE in kcal mol⁻¹) and binding constant (k^b in 88
- Table 5A.3 Binding energy (ΔE in kcal mol⁻¹) and binding constant (k^b in min⁻¹) of all the complexes with **HSA**, evaluated by molecular docking
- Table 5A.4 Hydrogen bond and metal-receptor interaction of 90 ruthenium(III) complexes with **HSA** evaluated by docking analysis
- Table 5A.5 Calculated bond length (Å) and bond angles (⁰) of 92 monoaquated and diaguated adduct
- Table 5A.6 Absolute energy values (in a.u.) of interacting adducts and calculated binding energy ($\Delta E \times 10^2$ in kcal mol⁻¹) of ruthenium complexes with **HSA** calculated by two layer ONIOM method in gas phase. Interacting part of HSA in aqueous phase is calculated by high level UB3LYP/(LANL2DZ+6-31G(d,p)) method
- Table 5B.1 Optimized Geometric parameters (bond length in Å and bond angles in ⁰) of all the stationary points in ligand exchange reaction of **Ha** with histidine and cysteine at the level of BLYP/DNP in gas phase
- Table 5B.2 Optimized Geometric parameters (bond length in Å and bond angles in ⁰) of all the stationary points in ligand exchange reaction of **IIb** with histidine and cysteine at the level of BLYP/DNP in gas phase
- Table 5B.3 Total relative energy (E_{tot}) , relative enthalpy (ΔH) and free energy (ΔG) values of **IIa** with histidine and cysteine calculated at BLYP/DNP level. Energy values are in kcal mol⁻¹
- Table 5B.4 Total relative energy (E_{tot}) , relative enthalpy (ΔH) and free energy (ΔG) values of diaqua ruthenium(III) complex **IIb** with His and Cys calculated at BLYP/DNP level. Energy values are in kcal mol⁻¹. Solvent phase values are in parenthesis

Rate constant values for the substitution reaction of IIa and Table 5B.5 114 **IIb** with histidine and cysteine Optimized geometric parameters (bond length in Å and bond Table 5C.1 121 angles in ⁰) of all the stationary points in chloride exchange reaction of Ia-HSA adduct with water at the level of B3LYP/(LANL2DZ+6-31G(d,p) in gas phase Table 5C.2 Optimized geometric parameters (bond length in Å and bond 124 angles in ⁰) of all the stationary points in water exchange reaction of Ia-HSA adduct with nitric oxide at the level of B3LYP/(LANL2DZ+6-31G(d,p) in gas phase Table 5C.3 Total ZPE $(E_{tot/ZPE})$, solvation energies (G_{solv}) , thermal 126 contributions to enthalpies (H_{therm}) , thermal contribution to Gibbs free energies (G_{therm}), change of enthalpies (ΔH) and change of Gibbs free energies (ΔG) 298.15 K for the species involved in hydrolysis process of Ia-HSA adduct. Energy values are in kcal mol⁻¹, entropy values are in cal k⁻¹ mol⁻¹ and rate constant value is in s⁻¹. Solvent phase values are in parenthesis Table 5C.4 128 Total ZPE $(E_{tot/ZPE})$, solvation energies (G_{solv}) , thermal contributions to enthalpies (H_{therm}) , thermal contribution to Gibbs free energies (G_{therm}), change of enthalpies (ΔH) and change of Gibbs free energies (ΔG) and rate constant at 298.15 K for the species involved in the nitrosylation reaction of hydrolyzed **Ia-HSA** adduct. Energy values are in kcal mol⁻¹, entropy values are in cal k⁻¹ mol⁻¹ and rate constant value is in s⁻¹. Solvent phase values are in parenthesis Table 6.1 Selected bond lengths (Å) and bond angles (⁰) calculated for 138 complex Ia and complex IVa at M062X / (LANL2DZ+6-31G (d,p)) level in the gas phase calculated bond lengths (Å) and bond angles (0) of ruthenium Table 6.2 141 complex-base pair adduct at M062X / (LANL2DZ+6-

31G(d,p)) level in the gas phase.

Table 6.3	Hydrogen bond distances of isolated base pair and base pair	144
	adducts calculated at M062X / (LANL2DZ+6-31G(d,p)) level	
Table 6.4	Absolute free energy values ($E_{rutehniumcomplex-basepair}$ in kcal mol ⁻¹)	147
	of interacting adducts and calculated interaction energies (ΔE	
	in kcal mol ⁻¹) calculated at M062X /(LANL2DZ+6-31G(d,p))	
	level of theory	
Table 6.5	Calculated dipole moment (μ in D) values of ruthenium	148
	complex-base pair adducts calculated at M062X	
	/(LANL2DZ+6-31G (d,p)) level	
Table 6.6	Natural atomic charges and natural electron configuration of	150
	the selected atoms ruthenium complex-base pair adducts and	
	isolated base pairs calculated at M062X /(LANL2DZ+6-31G	
	(d,p)) level	
Table 7.1	Bond distances (Å), bond angles (⁰) and dihedral angles (⁰) of	161
	the ruthenium(II) complexes and X-ray data for	
	$[Ru(dmp)_2(dppz)]^{2^+}$	
Table 7.2	Energies of HOMO (E_H in eV) and LUMO (E_L in eV) and	163
	chemical hardness (η in eV) of three ruthenium(II) complexes	
Table 7.3	Free energy of binding (ΔG in kcal mol ⁻¹), RMSD values and	163
	intrinsic binding constant (K _b in M ⁻¹) of docked structures	
Table 7.4	Hydrogen bond interaction of three ruthenium (II) complexes	164
	with d(ATATAT) ₂ and d(GCGCGC) ₂ sequences evaluated by	
	docking analysis	
Table 7.5	The structural parameters between of two AT and GC base	167
	pairs of DNA in free as well in complex forms	
Table 7.6	The calculated binding energy (ΔE in kcal mol ⁻¹) of I , II and	168
	III with d(ATATAT) ₂ and d(GCGCGC) ₂ duplex	

LIST OF FIGURES

Chapter 1 General Introduction

Fig. 1.1	Structures of cisplatin, carboplatin and oxaliplatin.	4
Fig. 1.2	Hydrolysis of cisplatin.	4
Fig. 1.3	Cisplatin form adducts with the N7 nitrogen of guanine	5
	bases (PDB code 3LPV).	
Fig. 1.4	The change of oxidation states of ruthenium in cancer and	8
	healthy cells.	
Fig. 1.5	Schematic representation of selective uptake of transferrin	9
	by cancer cells.	
Fig. 1.6	Ammine-chlorido derivatives.	9
Fig. 1.7	Structural formulas of dimethylsulfoxide complexes.	10
Fig. 1.8	Chemical structure of NAMI-A derivatives.	11
Fig. 1.9	Pyridine analogue of NAMI-A complex.	12
Fig. 1.10	Structural formulas of KP418 (Im[trans-RuCl ₄ (Him) ₂]),	12
	KP1019 (Ind[trans-RuCl ₄ (Hind) ₂]), KP1339 (Na[trans-	
	$RuCl4(Hind)_2]$).	
Fig. 1.11	Polypyridyl ligands.	13
Fig. 1.12	Structural formula of ruthenium Polypyridyl complexes.	13
Fig. 1.13	$[(\eta 6\text{-arene})Ru(II)(en)X]^+$ (where arene= benzene, p-	14
	cymene, biphenyl, 5,8,9,10-tetrahydroanthracene or 9,10-	
	dihydroanthracene, X is Cl or I) [($\eta 6$ -	
	arene) $Ru(II)(pta)XY$] (where R1, R2 = alkyl groups; X	
	and $Y = Cl/$ different μ -dicarboxylate ligands.	
Fig. 1.14	Schematic diagram of guanine and adenine.	17
	Chapter 2 Importance of computational chemistry	
Fig. 2.1	Small molecule fits to a large macromolecule	39
Chapter 3	Structure and reactivity of ruthenium(III) anticancer comple	exes
Fig. 3.1	Optimized geometries of ruthenium(III) complexes with	50
	appropriate numbering obtained from B3LYP/	
	(LanL2DZ+6-31G(d,p)) calculation.	
Fig.3.2	Plot of experimental and calculated cytotoxicity (log IC-1)	53

values of ruthenium complexes against HT-29 colon carcinoma cell line.

Chapter 4 hydrolysis mechanism of ruthenium(III) anticancer complexes

Fig. 4.1	Optimized structures for the species involved in the first	62
	step of Cl hydrolysis of complex II and IV (path 1).	
Fig. 4.2	Optimized structures for the species involved in the second	66
	step of Cl ⁻ hydrolysis of complex II and IV (path 2 and 3).	
Fig.4.3	Optimized structures for the species involved in the first	67-68
	DMSO hydrolysis of complex II and IV (path 4).	
Fig.4.4	Optimized structures for the species involved in the second	69
	DMSO hydrolysis of complex II and IV (path 5).	
Fig.4.5	Free energy profile diagram for the first hydrolysis of	72
	complex II (a) in gas phase and (b) in solvent phase.	
Fig.4.6	Free energy profile diagram for the first hydrolysis of	73
	complex IV (a) in gas phase and (b) in solvent phase.	
Fig.4.7	Free energy profiles of Cl ⁻ (path 1) and DMSO (path 4)	74
	hydrolysis of complexes II and IV.	
Fig.4.8	Free energy profile diagram for the second step of Cl	75
	(path 2) and DMSO (path 4) hydrolysis reaction of	
	complexes II and IV.	
Chapter	5A Interactions of the aquated forms of ruthenium(III) antica	ncer
	complexes with protein	
Fig.5A.1	Schematic 2D diagram of the model system for ruthenium	83
	complex bound to HSA binding site. Layers that are	
	partitioned are shown for ONIOM2 calculations. A is the	
	inner layer (QM calculations) and B is the outer layer	
	(MM calculations). The arrangement of the residues	
	shown in 2D diagram is not their actual position in 3D.	
Fig.5A.2	Optimized geometries of ruthenium(III) complexes with	86
	appropriate numbering obtained from B3LYP/	

	(LanL2DZ+6-31G(d,p)) calculation.	
Fig.5A.3	Docked structures of monoaquated adducts at the active	88
	site of protein receptor. The rest part of protein structure is	
	not shown for clarity.	
Fig.5A.4	Docked structures of diaquated adducts at the active site of	89
	protein receptor. The rest part of protein structure is not	
	shown for clarity.	
Fig.5A.5	Optimized geometries of monoaquated and diaquated	93
	adducts with appropriate numbering obtained from two	
	layer QM/MM method.	
Chapter 51	B Interaction mechanism of aquated form of ruthenium(III) ant	icancer
	complexes with histidine and cysteine	
Fig.5B.1	Optimized structures of R, TS, and P for interactions of	105
	Ha with histidine and cysteine calculated at BLYP/DNP	
	level.	
Fig.5B.2	Optimized structures of R, TS, and P for interactions of	108
	Hb complex with histidine and cysteine calculated at	
	BLYP/DNP level.	
Fig.5B.3	Free energy profile diagram of the monoaqua	109
	ruthenium(III) complex IIa binding to the histidine and	
	cysteine in the gas phase calculated at BLYP/DNP level.	
Fig.5B.4	Free energy profile diagram of monoaqua ruthenium(III)	110
	complex Ha binding to the histidine and cysteine in the	
	solvent phase calculated at BLYP/DNP level.	
Fig.5B.5	Free energy profile diagram of diaqua ruthenium(III)	113
	complex IIb binding to the histidine and cysteine in the	
	gas phase calculated at BLYP/DNP level.	
Fig.5B.6	Free energy profile diagram of diaqua ruthenium(III)	113
	complex IIb binding to the histidine and cysteine in the	
	solvent phase calculated at BLYP/DNP level.	

Chapter 5C Quantum chemical studies on detail mechanism of nitrosylation of ruthenium(III) complex-HSA adduct

Fig.5C.1	Optimized structures for the species involved in hydrolysis	122
	reaction of Ia-HSA adduct calculated at B3LYP/	
	(LanL2DZ and 6-31G(d,p)) level.	
Fig.5C.2	Optimized structures for the species involved in	125
	nitrosylation reaction of hydrolyzed Ia-HSA adduct	
	calculated at B3LYP/ (LanL2DZ and 6-31G(d,p)) level.	
Fig.5C.3	Free energy profile diagram of the hydrolysis step of Ia-	126
	HSA adduct in gas phase calculated at B3LYP/ (LanL2DZ	
	and $6-31G(d,p)$) level.	
Fig.5C.4	Free energy profile diagram of the hydrolysis step of Ia-	127
	HSA adduct in solvent phase calculated at B3LYP/	
	(LanL2DZ and 6-31G(d,p)) level.	
Fig.5C.5	Enthalpy profile diagram of the hydrolysis step of Ia-HSA	127
	adduct in solvent phase calculated at B3LYP/ (LanL2DZ	
	and $6-31G(d,p)$) level.	
Fig.5C.6	Free energy profile diagram of the nitrosylation step of Ia-	129
	HSA adduct in gas phase calculated at B3LYP/ (LanL2DZ	
	and 6-31G(d,p)) level.	
Fig.5C.7	Free energy profile diagram of the nitrosylation step of Ia-	129
	HSA adduct in solvent phase calculated at B3LYP/	
	(LanL2DZ and 6-31G(d,p)) level.	
Fig.5C.8	Enthalpy profile diagram of the nitrosylation step of	130
	NAMI-A-HSA adduct in solvent phase calculated at	
	B3LYP/ (LanL2DZ and 6-31G(d,p)) level.	
Chapter 6	Interaction of aquated form of ruthenium(III) anticancer con	mplexes
	with normal and mismatch base pairs	
Fig.6.1	Optimized geometries of complex Ia and complex IVa	139
	with appropriate numbering obtained from M062X /	
	(LanL2DZ+6-31G(d,p)) calculation.	
Fig.6.2	Optimized geometries of normal and mismatch base pair	139-140

	with appropriate numbering obtained from M062X/ (LanL2DZ+6-31G(d,p)) calculation.	
Fig.6.3	Optimized structures of interactions of complex Ia with	142
	normal and mismatch base pair obtained from M062X /	
	(LanL2DZ+6-31G(d,p)) calculation.	
Fig.6.4	Optimized structures of interactions of complex IVa with	142-143
	normal and mismatch base pair obtained from M062X /	
	(LanL2DZ+6-31G(d,p)) calculation.	
Fig.6.5	Frontier HOMO diagrams of normal (GC and AT) and	144-145
	mismatch base pair (GG, AA, TT and CC)	
Fig.6.6	Frontier LUMO diagrams of complex Ia and IVa.	145
Fig.6.7	Base pair-ruthenium complex adduct (HOMO of base pair	146
	and LUMO of ruthenium complex).	
Chapter 7 In	teraction of ruthenium(II) antitumor complexes with d(ATAT	$(AT)_2$ and
	$d(GCGCGC)_2$	
Fig.7.1	Docked conformation of native ligand [Ru(phen) ₂ dppz] ²⁺	159
	as compared to the conformation of ligand in the original	
	crystal structure.	
Fig.7.2	Optimized geometries of ruthenium(II) complexes with	162
	appropriate numbering obtained from B3LYP/	
	(LanL2DZ+6-311+G(d,p)) calculation.	
Fig.7.3	Docked conformation of complex I with $d(ATATAT)_2$ and	165
	$d(GCGCGC)_2$ sequence.	
Fig.7.4	Optimized geometry of d(ATATAT) ₂ and d(GCGCGC) ₂	166
	obtained by two layer QM/MM method.	
Fig.7.5	Optimized geometries of d(ATATAT) ₂ —I and	167
	d(GCGCGC) ₂ —I adducts obtained by two layer QM/MM	
	method.	

CONTENTS

Acknowline	wledgement	1-11
 Abstrac 	et	iii-viii
 List of Abbreviations 		ix-x
List of	List of Tables	
List of	Figures	xv-xix
	Chapter 1 General Introduction	1-27
1.1	Cancer	1-3
1.1.1	Types of cancer	1-2
1.1.2	Different existing cancer treatments	2-3
1.2	Cancer treatments by Platinum Therapies	3-6
1.2.1	Bases of modern chemotherapy	3-4
1.2.2	Mode of action	4-5
1.2.3	From platinum to ruthenium	5-6
1.3	Ruthenium anticancer complexes	6-16
1.3.1	Ruthenium properties that make it suitable for biological	1 7-9
	applications	
1.3.2	Classification of ruthenium complexes with anticancer	r 9-14
	properties	
1.3.2.1	Ammine-chlorido derivatives	9
1.3.2.2	Dimethyl sulfoxide complexes	10
1.3.2.3	NAMI-A type of complexes	10-11
1.3.2.4	Complexes with other heterocyclic ligands	11-12
1.3.2.5	Ruthenium polypyridyl complexes	12-13
1.3.2.6	Organoruthenium complexes	13-14
1.3.3	Ruthenium complexes undergoing clinical trials	14-16
1.3.3.1	NAMI-A	14-15
1.3.3.2	KP1019	16
1.3.4	Biological target of ruthenium drug	16
1.4	Computational approach	17-18
1.5	Scope of the work	18-19
		20-27

	Chapter 2 Importance of computational chemistry	28-44
2.1	Computational chemistry	28
2.1.1	Molecular mechanics	28
2.1.2	Quantum mechanics	28
2.1.2.1	The Schrödinger equation	28-29
2.1.2.2	Born-Oppenheimer approximation	29-30
2.1.2.3	Hartree-Fock method	30-31
2.1.2.4	Density functional theory	31-35
2.1.2.5	DFT based reactivity descriptors	35-37
2.1.3	Quantum mechanics/molecular mechanics methods(QM/MM)	37-38
2.1.4	Molecular docking	38-39
2.1.4.1	Mechanics of docking	39-40
2.1.4.2	Application of molecular docking	40
2.1.4.3	Summary:	40-41
	References	42-44
Chaj	oter 3 Structure and reactivity of ruthenium(III) anticancer	45-57
	complexes	
3.1	Introduction	45-46
3.2	Computational details	47
3.3	Results and discussion	47-53
3.3.1	Structure	47-50
3.3.2	Global reactivity	50-51
3.3.3	Local reactivity	51-52
3.3.4	QSAR study	52-53
3.4	Conclusion	53-54
	References	55-57
Cha	pter 4 Hydrolysis mechanism of ruthenium(III) anticancer	58-79
	complexes	
4.1.	Introduction	58-59
4.2	Computational Details	59-60
4.3	Results and Discussion	60-75

4.3.1	Structural characteristics of the species involved in Cl-hydrolysis	60-66
4.3.1.1	First hydrolysis step	61-62
4.3.1.2	Second step of hydrolysis	63-66
4.3.2	Structural characteristics for the species involved in DMSO	66-69
1.3.2	hydrolysis	00 07
4.3.3	Reaction Profiles of the Hydrolysis Processes	69-75
4.3.3.1	Comparison of first and second step of Cl ⁻ hydrolysis	69-73
4.3.3.2	Comparison of first step of Cl and first step of DMSO	73-74
4222	hydrolysis	74.75
4.3.3.3	Comparison of second step of Cl ⁻ and DMSO hydrolysis	74-75
4.4	Conclusion	76
	References	77-79
Chapt	er 5A Interactions of the aquated forms of ruthenium(III)	80-99
-	anticancer complexes with protein	
5A.1	Introduction	80-81
5A.2.1	Computational Detail	82
5A.2.2	Molecular docking simulation	82-83
5A.2.3	QM/MM calculation	83-84
5A.3	Results and Discussion	84-95
5A.3.1	Structural analysis of monoaqua and diaqua complexes	84-86
5A.3.2	Stability of the ruthenium complexes	87
5A.3.3	Docking study	87-89
5A.3.4	ONIOM study	90-95
5A.3.4.1	Structural Characteristics	90-93
5A.3.4.2	Stability	93-95
5A.4	Conclusion	95-96
	References	97-99
Chapter 3	5B Interaction mechanism of aquated form of ruthenium(III)	100-116
	anticancer complexes with histidine and cysteine	
5B.1	Introduction	100-101
5B.2	Computational Methods	102-103

5B.3	Results and Discussion	103-114
5B.3.1	Structural Characteristics	103-108
5B.3.2	Energy Profiles of ligand exchange reaction	108-113
5B.3.3	Kinetic analysis	113-114
5B.4	Conclusion	114
	References	115-116
Cha	pter 5C Quantum chemical studies on detail mechanism of	117-134
	nitrosylation of ruthenium(III) complex-HSA adduct	
5C.1	Introduction	117-118
5C.2	Computational Details	118-119
5C.3	Results and Discussion	119-130
5C.3.1	Structural characteristics	119-125
5C.3.2	Energy profiles	125-130
5C.3.3	Kinetic analysis	130
5C.4	Conclusion	130-131
	References	132-134
Chapte	r 6 Interaction of aquated form of ruthenium(III) anticancer	135-154
	complexes with normal and mismatch base pairs	
6.1	Introduction	135-136
6.2	Computation details	137
6.3	Results and Discussion	137-150
6.3.1	Structural analysis	137-140
6.3.2	Structural characteristics of ruthenium complexes-base pair adduct	140-143
6.3.3	Interaction in ruthenium complex -base pair adduct	143-146
6.3.4	Stability of ruthenium complex-base pair adduct	146-148
6.3.5	Natural Bond orbital (NBO) analysis	149-150
6.4	Conclusion	151
	References	152-154
Chap	ter7 Interaction of ruthenium(II) antitumor complexes with	155-173
	$d(ATATAT)_2$ and $d(GCGCGC)_2$	
7.1	Introduction	155-157

	Chapter 8 Conclusion	174-177
	References	170-173
7.4	Conclusion	169
7.3.4	QM/MM study	165-168
7.3.3	Molecular docking study	163-165
7.3.2	Stability of the ruthenium complexes	162-163
7.3.1	Structural analysis of metal complexes	160-162
7.3	Results and Discussion	160-168
7.2.3	Ruthenium complex-DNA interaction by QM/MM method	159-160
7.2.2	Automated DNA— ruthenium complex docking	158-159
7.2.1	Structure	157
7.2	Computational Details	157-160

Appendix I : List of conferences and workshops attended

Appendix II : Reprints of published papers