DECLARATION

I, Nivedita Dasgupta bearing Registration No. Ph.D/2415/13 Dt. 12.09.2013, hereby declare that the subject matter of the thesis entitled "Molecular Characterization, Gene Location and Transferability of Quinolone Resistance among Enterobacteriaceae" is the record of work done by me and the contents of this thesis did not form the basis for award of any degree to me or to anybody else to the best of my knowledge. The thesis has not been submitted in any other University/ Institute. Also, it is declared that the content of the thesis is free of plagiarism.

Place: (Nivedita Dasgupta)

Date: Registration No. Ph.D/ 2415/13 Dt. 12.09.2013

Acknowledgements

I extend my deepest and sincere gratitude to my supervisor Dr. Amitabha Bhattacharjee, Assitant Professor, Dept. of Microbiology, Assam University, Silchar; for his guidance and his inspiration. I greatly appreciate his persistent care with positive suggestions in all significant circumstances during the entire period of my research work. I want to thank him for his unstinting supports which enable me to accomplish the research work successfully.

I would like express my utmost thanks to my co-supervisor Dr. Debadatta Dhar (Chanda), Associate Professor, Dept. of Microbiology, Silchar Medical College and Hospital, Silchar; for her help during my research work.

I express my profound gratitude to Prof. Piyush Pandey, HOD, Department of Microbiology, Assam University, Silchar for his able guidance and kind support in each and every step of my research work. I am also thankful to him for his constant encouragement in completion of this work.

I convey my sincere gratitude to Prof. Sankar Kumar Ghosh, Dean, Hargobind Khurana School of Life Sciences and Dr. Indu Sharma, an Assistant Professor, Department of Microbiology, Assam University, Silchar; for their encouragement and valuable suggestions in my research work.

Also take this opportunity to thank the Technical, Laboratory and Office Staffs of the department for their service and assistance during the process. I am grateful to all the people I have along the way and have contributed to the development of research work. I acknowledge my sincere thanks to Mr. Deepjyoti paul and Mr. Birson Ingti and to entire team of research scholars of ARRL for technical support and cooperation during my research work.

I am deeply thankful to my family members specially my parents (Late. Nihar Ranjan Dasgupta and Mrs. Nita Dasgupta) and my son (Anubhav DattaChoudhury) for their love, support, sarcrifice, untiring efforts and countless prayers. I want to dedicate this

thesis to the loving memory of my father whose role in my life was and remains immense. The last word of acknowledgement for my dear husband, Dr. Arunooday DattaChoudhury, without whom this thesis would have not been possible. I want to thank him for having faith in me and for being the greatest support during my tough days.

Last but not the least; I bow my head to Almighty, who has blessed upon me the strength to accomplish the piece of task successfully.

(Nivedita Dasgupta)

LIST OF CONTENTS

Contents	Page no.
1. Introduction	1-7
2. Review of literature	8- 64
2.1 Historical Background	8
2.1.1 Generation and development of quinolones	9-16
2.2 Mechanisms of quinolone action	16
2.2.1 DNA gyrase	18
2.2.2 Topoisomerase IV 2.2.2.1 Mode of action	20 20
	21
2.3 Quinolone resistance- mechanism of action:	21
2.3.1 Target alterations	23
2.3.1.1 Alterations in the DNA gyrase 2.3.1.2 Alteration in topoisomerase IV	25
2.3.2 Decreased uptake	26
2.3.3 Transferability of quinolone resistance	29
2.3.3.1 Pentapeptide repeat proteins	30-33
2.3.3.1.1 About <i>Qnr</i> proteins	33
2.3.3.2 Chromosomal <i>qnr</i> Genes	34
2.3.3.3 Mechanism of action of pentapeptide	
repeats protein families conferring quinolone resistance	36
2.3.3.4 Origins of <i>qnr</i> genes	38-39
2.3.3.5 Biological functions of <i>qnr</i>	38
2.3.3.6 Acc(6')-Ib-cr	40
2.3.3.6.1 Resistance Activity of AAC(6')-	41
<i>Ib-cr</i>	
2.3.3.6.2 AAC(6')-Ib-cr Protein	41
2.3.3.3.6.3 Genetic Environment of	42
acc(6')-Ib-cr Plasmids	
2.3.4 Plasmid-Mediated Quinolone Efflux	43
2.3.4.1 Genetic Environment of Efflux-Encoding Plasmids	44
2.4 Status of research related to quinolone resistance	45
2.4.1 Status in India	51
2.5 Methods of Detection of PMQR	52
2.5.1 Molecular detection of quinolone resistance	53
2.6 Genetic environment of <i>qnr</i> determinants	59
2.7 Therapeutic option	64
3. Materials and methods	65-119
3.1 Samples:	65
3.1.1 Patient population	65
3.1.2 Environmental samples	65
3.2 Isolates for the study	66
3.3 Isolation and identification of isolates from clinical	67
specimens	
3.3.1 Gram staining	67
3.3.2 Motility (hanging drop preparation)	68
3.3.3 Biochemical test	69

3.3.3.1 Catalase test	69
3.3.3.2 Oxidase test	69
3.3.3.3 Indole test	70
3.3.3.4 Methyl red test	71
3.3.3.5 Voges proskauer test	71
3.3.3.6 Simmons citrate reactions	72
3.3.3.7 Urease test	73
3.3.3.8 Triple sugar iron test	73
3.3.3.9 Phenyl alanine deaminase test	75
3.3.3.10 Amino acid decarboxylase	75
and dehydrolase test (Moller's method)	73
3.3.3.11 Sugar fermentation	77
-	78
3.3.3.12 Oxidative-Fermentation test (O/F test/	/ 6
Hugh and Leifson test)	70
3.3.3.13 Nitrate reduction test	79
3.4 Antibiotic susceptibility testing	79
3.4.1 Disc Diffusion Method	79
3.4.1.1 Procedure for the Preparation of a 0.5	81
McFarland Standard	
3.4.2 Minimum inhibitory concentration (MIC) study	83
3.4.2.1 Preparation of stock solution	83
3.4.2.2 Preparation of agar dilution plates	83 83
3.4.2.3 Preparation of agar plates	83
3.4.2.4 Preparation of inoculum	
3.4.2.5 Inoculation of test isolates in antibiotic	83
agar plates	84
3.4.2.6 Incubation of antibiotic agar plate	
3.4.2.7 Preparation of the stock solution	85
3.5 Genotypic characterization of <i>qnr</i>	85
3.5.1 Multiplex PCR1	86
3.5.1.1 Preparation of DNA template	86
3.5.1.2 Boiling centrifugation method	86
3.5.1.3 Preparation of reaction mixture	86
3.5.1.4 Reaction condition	86
3.5.1.5 Gel electrophoresis and analysis of PCR	95
products	0.0
3.5.2 Multiplex PCR2	88
3.5.2.1 Preparation of DNA template	88 88
3.5.2.2 Preparation of reaction mixture	88
3.5.2.3 Reaction condition	89
3.5.2.4 Gel electrophoresis and analysis of PCR	89
products	00
3.5.2.5 Preparation of TBE (10X)	89
3.6 PCR product purification	90
3.7 Sequencing of all <i>qnr</i> genes	91
3.8 Transferability of <i>qnr</i> genes	91
3.8.1 Isolation of plasmid	91
3.8.2 Transformation of plasmid and their role in	93

antibiotic resistance in recipient	
3.8.2.1 Preparation of competent cells	93
3.8.2.2 Transformation procedure	95
3.8.2.3 Component of Luria Bertani agar	96
(LB agar) (Per litre)	07
3.8.2.4 Preparation of stock solution	97
3.8.2.5 Composition of SOC medium (per litre)	97
3.8.2.6 Preparation of LB norfloxacin, LB	97
ciprofloxacin, LB ofloxacin, LB levofloxacin agar plates	
3.8.2.7 Spreading of cells on LB agar plates	97
3.8.2.8 PCR detection of <i>qnr</i> gene in transformed	98
recipient	76
3.8.2.9 Minimum inhibitory concentration of	98
transformants	
3.8.3 PCR detection of Incompatibility group (Inc)	98
in transformants	
3.8.3.1 Targeted gene in multiplex PCR	98
3.8.3.2 Targeted gene in simplex PCR	98
3.8.3.3 Preparation of DNA template	
3.8.3.4 Primers	99
3.8.3.5 Preparation of reaction mixture	99
3.8.3.6 Reaction condition for multiplex PCR	99
3.8.3.7 Reaction condition for simplex PCR	99
3.8.3.8 Visualization of PCR products	99
3.8.4 Transfer of resistance determinants	101
through conjugation	101
3.8.4.1 Strain description	101
3.8.4.2 Donor strain	102
3.8.4.3 Strain B (Recipient strain)	102
3.8.4.4 Preparation of antibiotic stock solution	102
3.8.4.5 Preparation of Streptomycin stock	102
3.8.4.6 Revival of donor strain and strain B	102
3.8.4.7 Preparation of bacteria	103
3.8.4.8 Procedure of conjugation	103
3.8.4.9 PCR detection of <i>qnr</i> gene in	103
transconjugants	104
3.8.4.10 PCR detection of Incompatibility group	104
(Inc) in transconjugants	101
3.8.3.11 Minimum inhibitory concentration of	104
transconjugants	
3.9 DNA fingerprinting of quinolone resistant determinants	105
3.9.1 Enterobacterial repititive intergenic	
consensus (ERIC) sequences PCR	105
3.9.1.1 Preparation of DNA template	105
3.9.1.2 Preparation of reaction mixture	105
3.9.1.3 Reaction condition	105
3.9.1.4 Visualization of PCR products	105
•	103

3.10 Determination of genetic environment 3.10.1 Detection of variable region of integron among quinolone resistant isolates 3.10.1.1 Preparation of DNA template 3.10.1.2 Primers	106 106 106 106
quinolone resistant isolates 3.10.1.1 Preparation of DNA template 3.10.1.2 Primers	106 106
quinolone resistant isolates 3.10.1.1 Preparation of DNA template 3.10.1.2 Primers	106
3.10.1.1 Preparation of DNA template 3.10.1.2 Primers	106
3.10.1.2 Primers	
	106
3.10.1.3 Preparation of reaction mixture	106
3.10.1.4 Reaction condition	106
3.10.1.5 Electrophoresis and Visualization of PCR	
products	107
^	10,
3.10.2 Detection of the location of qnr and $aac(6')$ -Ib-	107
cr gene within integron region	107
3.10.2.1 Preparation of DNA template	107
3.10.2.2 Primers	107
3.10.2.3 Preparation of reaction mixture	108
3.10.2.4 Reaction condition	
3.10.2.5 Electrophoresis and Visualization of PCR	108
products	100
3.10.2.6 Gel extraction and purification of	109
amplified PCR product	100
3.10.2.7 Sequencing	109
3.11 Association of <i>qnr</i> and <i>aac</i> (6') <i>Ib-cr</i> genes with mobile	109
element	110
3.11.1 Preparation of DNA template	
3.11.2 Preparation of reaction mixture	110
3.11.3 Reaction condition	110
3.11.4 Electrophoresis and Visualization of PCR products	110
3.11.5 Gel extraction and purification of amplified PCR	
product	111
3.11.6 Sequencing	111
3.12 Study of mutation in the quinolone resistant determining	
region by denaturing gradient gel electrophoresis	111
3.12.1 Preparation of DNA template	111
3.12.2 Primers	111
3.12.3 Preparation of reaction mixture	112
3.12.4 Reaction condition	112
3.12.5 Gel electrophoresis and analysis of PCR products	113
3.12.6 Denaturing Gradient Gel Electrophoresis (DGGE)	113
3.12.6.1 Reagent preparation	114
3.12.6.2 Sample Preparation	115
3.12.6.3 Temperature Controller	116
3.12.6.3.1 Pre-heating the running buffer	116
3.12.6.4 Assembling the Parallel Gradient Gel	116
Sandwich]
3.12.6.5 Casting parallel Denaturing Gradient	116
Gels	
3.12.6.6 Electrophoresis	117
3.12.6.6.1 Assembling the upper buffer	118
chamber	

3.12.6.6.2 Sample loading	118
3.12.6.6.3 Running the Gel	118
3.12.6.7 Staining and photographing the gel	118
3.12.6.8 Gel extraction and purification of	119
amplified PCR product	119
3.12.6.9 Sequencing	110
1 0	119
4. Results	131-190
4.1 Isolation and identification of isolates from clinical	120
specimens and environmental samples	120
4.1.1 Identification of isolates by biochemical test	121
4.2 Samples	123
4.2.1 Clinical isolates	123
4.2.2 Environmental isolates	125
4.3 Phenotypic screening of quinolone resistance	125
4.3.1: Resistance tree based on quinolone antibiotic	120
resistance pattern	129
4.3.2 Antibiotic susceptibility of environmental isolates	130
against quinolone antibiotics	130
4.4 Minimum inhibitory concentration	131
4.4 Willimitan numberly concentration	151
4.4.1 Minimum inhibitory concentration against	131
quinolones in clinical isolates	151
4.4.2 Minimum inhibitory concentration against	138
quinolones in environmental isolates	
4.5 Genotypic characterization of quinolone resistant isolates	141
4.5.1 Genotypic characterization of quinolone resistant	141
clinical isolates	
4.5.2 Genotypic characterization of quinolone resistant	
environmental isolates	141
4.6 Sequencing of all <i>qnr</i> genes and <i>aac</i> (6') <i>Ib-cr</i>	146
4.7 Transferability of <i>qnr</i> genes and <i>aac</i> (6') <i>Ib-cr</i> genes:	151
4.7.1 PCR detection of <i>qnr</i> genes and <i>aac</i> (6') <i>lb-cr</i> genes	
in transformants	151
	151
4.7.2 Selection specificity of <i>qnr</i> genes	152
4.7.3 MIC of transformants	155
4.8 Analysis of plasmids	155
4.9 Plasmid Incompatibility typing	164
4.10 Analysis of conjugative transferability of plasmid	164
4.10.1 PCR detection of <i>qnr</i> among transconjugants	
4.10.2 MIC of transconjugants	164
4.11 DNA fingerprinting of quinolone resistant isolates by ERIC	169
PCR	1.60
4.12 Determination of genetic environment	169
4.12.1 Detection of the location of <i>qnr</i> gene within	169
integron region	
4.12.2 Association of <i>qnr</i> and <i>aac</i> (6') <i>Ib-cr</i> genes with	169
mobile element	109
4.13 Study of mutation in the quinolone resistant determining	171
1 1	±/±

region by denaturing gradient gel electrophoresis:	
4.13.1 Sequencing of gyrA and parC genes	171
4.14 Therapeutic option	177
	150 105
5. Discussion	179-195
6. Conclusion	196-198
7. References	199-238
8. Appendices	
a. Abbreviations	i-ii
b. Composition of media	iii-ix
c. Publication	X

List of Figures

Fi	gure		Page
Nu	mber	Description	Number
	1.	Common structure of 4- quinolones	10
	2.	Structural development of 7- piperazinylquinolone from primary quinolones	12
	3.	Quinolone containing pyrrolidine skeleton at C-7 position	14
	4.	Quinolone with modification at C-7 and N-1 position	15
	5.	Mechanism of action of quinolone antibiotic	17
	6.	Generalized structure of the integron/gene cassette system	59
	7.	Genetic environment of plasmid mediated <i>qnr</i> A, <i>qnr</i> B and <i>qnr</i> S alleles	61
	8.	Schematic presentation of the genetic environment of <i>qnr</i> B19 genes	63
	a.	Colony morphology of <i>E. coli</i> on macconkey agar	124
9.	b.	Colony morphology of <i>Klebsiella</i> on macconkey agar	124
	C.	Colony morphology of <i>Proteus</i> on macconkey agar	124
	10.	Antibiotic susceptibility plates against quinolone antibiotics	126
	11.	Proportion of quinolone resistant enterobacterial hospital (n=79) and community (n=130) isolates at species level	128
	12.	Minimum inhibitory concentration of quinolone resistant isolates against ofloxacin at 2-256µg/ml	137
	13.	PCR amplification of <i>qnr</i> and <i>aac(6')-Ib-cr</i> genes in clinical isolates	144

		PCR amplification of <i>qnr</i> and <i>aac(6')-Ib-cr</i> genes	
14.		in environmental isolates	145
15.		Electropherogram of <i>qnr</i> A1 PCR amplicon sequence	146
16.		Electropherogram of <i>qnr</i> B7 PCR amplicon sequence	147
17.		Electropherogram of <i>qnr</i> S1PCR amplicon sequence	148
18.		Electropherogram of <i>qnr</i> D1PCR amplicon sequence	149
19.		Electropherogram of <i>aac(6')-Ib-cr</i> variants PCR amplicon sequence	150
20.		Transformants in <i>E. coli</i> JM107 recepient strain selected against quinolone antibiotics	152
21.		Analysis of plasmid of transformants harbouring qnr genes	156
22.		PCR detection of Inc groups in transformants	157
23.	Transconjugants in <i>E.coli</i> F strain (small white colonies)		165
	a.	Antibiotic susceptibility of transformant against quinolone antibiotics	166
24.	b.	Antibiotic susceptibility of <i>E.coli</i> JM 107 without plasmid against quinolone antibiotics	166
	c.	MIC of transformants against quinolone antibiotic	166
		MIC of transconjugants against quinolone	
	d.	antibiotic	166
25.	a.	DNA finger printing of <i>E.coli</i> by ERIC PCR	170
	_	DNA finger printing of <i>Klebsiella pneumonia</i> and	
	b.	Klebsiella oxytoca	170
	c.	DNA finger printing of <i>Proteus vulgaris</i> and <i>Proteus mirabilis</i>	170

26.		PCR amplification of <i>gyr</i> A (586bp) and <i>par</i> C (265bp)	172
27.	a.	Denaturing gradient gel electrophoresis pattern of gyrA gene	173
	b.		173
28.	a.	Sequence alignment of the four type of mutational pattern <i>gyr</i> A sequences with the sequence of <i>Escherichia coli</i> strain 493/89(EC493/89)	174
28.	b.	Sequence alignment of the four type of mutational pattern <i>par</i> C sequences with the sequence of <i>Escherichia coli</i> strain 493/89(EC493/89)	175
29.	a.	Protein sequence alignment of four mutational patterns of GyrA with the sequence of <i>Escherichia coli</i> strain 93/89 (EC493/89)	176
29.	b.	Protein sequence alignment of four mutational patterns of ParC with the sequence of <i>Escherichia coli</i> strain 493/89(EC493/89)	176
30.		Antibiotic susceptibility of quinolone resistant isolates	177

List of tables

Table		Page
Number	Description	Number
1.	Clinical development of quinolone Drug	13
2.	Comparison of quinolone Generations	16
3.	Mutation described in subunit <i>ParC</i> and <i>ParE</i>	22
4.	Mutation described in subunits <i>GyrA</i> and <i>GyrB</i> of resistant strain of <i>E. coli</i>	23
5.	Types of clinical specimen obtained for the study	66
6.	Summary of TSI reactions	74
7.	Antimicrobial agents used for the study	82
8.	Preparation of plates for MIC	85
9.	Oligonucleotide used as primers for amplification of different <i>qnr</i> genes	87
10.	Composition of buffer for isolation of plasmids	92
11.	Primers used for PCR based replicon typing	100-101
12.	Primers used for detection of variable region of integron	106
13.	Primers used for the detection of gene cassette within integron region	107
14.	Primers used for the association of <i>qnr</i> genes with mobile element	110

		1
15.	Primers used for the amplification of quinolone resistant determining region	112
16.	Cultural characteristics	121
17.	List of biochemical test	122
18.	Different isolates obtained from clinical sample in the study	123
19.	Different isolates obtained environmental sample in the study	125
20.	Phenotypic screening results of quinolone resistance among clinical isolates	127
21.	Antibiogram pattern (tree) of quinolone resistant clinical isolates	129
22.	Phenotypic screening results of quinolone resistance among environmental isolates	130
23.	MIC of the screened clinical isolates against norfloxacin	132
24.	MIC of the screened clinical isolates against ciprofloxacin	133
25.	MIC of the screened clinical isolates against ofloxacin	134
26.	MIC of the screened clinical isolates against gatifloxacin	135
27.	MIC of the screened clinical isolates against levofloxacin	136
28.	MIC of screened environmental isolates against norfloxacin	138
29.	MIC of screened environmental isolates against ciprofloxacin	139

30.	MIC of screened environmental isolates against ofloxacin	139
31.	MIC of screened environmental isolates against levofloxacin	140
32.	Distribution of <i>qnr</i> genes among test clinical isolates (harbouring single <i>qnr</i> gene)	142
33.	Distribution of <i>qnr</i> genes among test clinical isolates (harbouring multiple <i>qnr</i> gene)	143
34.	Distribution of <i>qnr</i> genes among test environmental isolates	145
35.	MIC ₅₀ and MIC ₉₀ of transformants harbouring qnrA	153
36.	MIC ₅₀ and MIC ₉₀ of transformants harbouring qnrB	153
37.	MIC ₅₀ and MIC ₉₀ of transformants harbouring <i>qnr</i> D	154
38.	MIC ₅₀ and MIC ₉₀ of transformants harbouring qnrS	154
39.	MIC ₅₀ and MIC ₉₀ of transformants harbouring $aac(6')Ib$ -cr	155
40.	Incompatibility typing of transformant harbouring <i>qnr</i> A	158
41.	Incompatibility typing of transformant harbouring <i>qnr</i> B	159

42.	Incompatibility typing of transformant harbouring <i>qnr</i> D	160
43.	Incompatibility typing of transformant harbouring <i>qnr</i> S	161
44.	Incompatibility typing of transformant harbouring <i>aac</i> (6') <i>Ib-cr</i>	162
45.	Incompatibility typing of transformant harbouring multiple <i>qnr</i> genes	163
46.	MIC ₅₀ and MIC ₉₀ of transconjugants harbouring <i>qnr</i> A	164
47.	MIC ₅₀ and MIC ₉₀ of transconjugants harbouring <i>qnr</i> B	167
48.	MIC ₅₀ and MIC ₉₀ of transconjugants harbouring <i>qnr</i> D	167
49.	MIC ₅₀ and MIC ₉₀ of transconjugants harbouring <i>qnr</i> S	168
50.	MIC ₅₀ and MIC ₉₀ of transconjugants harbouring <i>aac</i> (6') <i>Ib-cr</i>	168
51.	Antibiogram profile of quinolone resistant isolates	190