

CHAPTER VII: REFERENCES

- Adrian, N. R., Robinson, J. A., and Suflita, J. M. (1994). Spatial variability in biodegradation rates as evidenced by methane production from an aquifer. *Appl Environ Microbiol* *60*, 3632-3639.
- Allain, C. C., Henson, C. P., Nadel, M. K., and Knoblesdorff, A. J. (1973). Rapid single-step kinetic colorimetric assay for lactate dehydrogenase in serum. *Clinical chemistry* *19*, 223-227.
- Altschul, S. F., Gish, W., Miller, W., Myers, E. W., and Lipman, D. J. (1990). Basic local alignment search tool. *Journal of molecular biology* *215*, 403-410.
- Alvarado, A., Montañez-Hernández, L. E., Palacio-Molina, S. L., Oropeza-Navarro, R., Luévanos-Escareño, M. P., and Balagurusamy, N. (2014). Microbial trophic interactions and *mcrA* gene expression in monitoring of anaerobic digesters. *Frontiers in Microbiology* *5*.
- Amann, R. I., Ludwig, W., and Schleifer, K.-H. (1995). Phylogenetic identification and in situ detection of individual microbial cells without cultivation. *Microbiological reviews* *59*, 143-169.
- An, Z., Xie, L., Cheng, H., Zhou, Y., Zhang, Q., He, X., and Huang, H. (2009). A silver staining procedure for nucleic acids in polyacrylamide gels without fixation and pretreatment. *Analytical biochemistry* *391*, 77-79.
- Anderson, I., Wirth, R., Lucas, S., Copeland, A., Lapidus, A., Cheng, J.-F., Goodwin, L., Pitluck, S., Davenport, K., and Detter, J. C. (2011). Complete genome sequence of *Staphylothermus hellenicus* P8T. *Standards in genomic sciences* *5*, 12.
- Angel, R., Claus, P., and Conrad, R. (2012). Methanogenic archaea are globally ubiquitous in aerated soils and become active under wet anoxic conditions. *The ISME journal* *6*, 847-862.

- Antony, C. P., Murrell, J. C., and Shouche, Y. S. (2012). Molecular diversity of methanogens and identification of *Methanobrevibacter* sp. as active methylophilic Archaea in Lonar crater lake sediments. *FEMS microbiology ecology* *81*, 43-51.
- Aslanian, A. (2006). Monitoring of bacterial community changes in leachate and wetland soil using T-RFLP analysis. *International journal of environmental studies* *63*, 143-153.
- Bapteste, É., Brochier, C., and Boucher, Y. (2005). Higher-level classification of the Archaea: evolution of methanogenesis and methanogens. *Archaea* *1*, 353-363.
- Bardgett, R. D., and van der Putten, W. H. (2014). Belowground biodiversity and ecosystem functioning. *Nature* *515*, 505-511.
- Barns, S. M., Delwiche, C. F., Palmer, J. D., and Pace, N. R. (1996). Perspectives on archaeal diversity, thermophily and monophyly from environmental rRNA sequences. *Proceedings of the National Academy of Sciences* *93*, 9188-9193.
- Bassi, N., Kumar, M. D., Sharma, A., and Pardha-Saradhi, P. (2014). Status of wetlands in India: a review of extent, ecosystem benefits, threats and management strategies. *Journal of Hydrology: Regional Studies* *2*, 1-19.
- Bloom, A. A., Palmer, P. I., Fraser, A., Reay, D. S., and Frankenberg, C. (2010). Large-scale controls of methanogenesis inferred from methane and gravity spaceborne data. *Science* *327*, 322-325.
- Bintrim, S. B., Donohue, T. J., Handelsman, J., Roberts, G. P., and Goodman, R. M. (1997). Molecular phylogeny of Archaea from soil. *Proceedings of the National Academy of Sciences* *94*, 277-282.
- Bogard, M. J., del Giorgio, P. A., Boutet, L., Chaves, M. C. G., Prairie, Y. T., Merante, A., and Derry, A. M. (2014). Oxidic water column methanogenesis as a major component of aquatic CH₄ fluxes. *Nat Commun* *5*.

- Bogner, J., Pipatti, R., Hashimoto, S., Diaz, C., Mareckova, K., Diaz, L., Kjeldsen, P., Monni, S., Faaij, A., and Gao, Q. (2008). Mitigation of global greenhouse gas emissions from waste: conclusions and strategies from the Intergovernmental Panel on Climate Change (IPCC) Fourth Assessment Report. Working Group III (Mitigation). *Waste Management & Research* 26, 11-32.
- Borrel, G., Harris, H. M., Tottey, W., Mihajlovski, A., Parisot, N., Peyretailade, E., Peyret, P., Gribaldo, S., O'Toole, P. W., and Brugère, J.-F. (2012). Genome sequence of “*Candidatus Methanomethylophilus alvus*” Mx1201, a methanogenic archaeon from the human gut belonging to a seventh order of Methanogens. *Journal of bacteriology* 194, 6944-6945.
- Borrel, G., O'Toole, P. W., Harris, H. M., Peyret, P., Brugère, J.-F., and Gribaldo, S. (2013). Phylogenomic data support a seventh order of methylotrophic methanogens and provide insights into the evolution of methanogenesis. *Genome biology and evolution* 5, 1769-1780.
- Brablcová, L., Buriánková, I., Badurová, P., Chaudhary, P. P., and Rulík, M. (2014). Methanogenic archaea diversity in hyporheic sediments of a small lowland stream. *Anaerobe*.
- Brauer, S. (2012). Seasonal changes in methanogenesis and methanogenic community in three peatlands, New York State.
- Bridgham, S. D., Cadillo-Quiroz, H., Keller, J. K., and Zhuang, Q. (2013). Methane emissions from wetlands: biogeochemical, microbial, and modeling perspectives from local to global scales. *Global Change Biology* 19, 1325-1346.
- Brochier-Armanet, C., Gribaldo, S., and Forterre, P. (2012). Spotlight on the Thaumarchaeota. *ISME Journal-International Society for Microbial Ecology* 6, 227.
- Cadillo-Quiroz, H., Bräuer, S., Yashiro, E., Sun, C., Yavitt, J., and Zinder, S. (2006). Vertical profiles of methanogenesis and methanogens in two contrasting acidic

- peatlands in central New York State, USA. *Environmental Microbiology* 8, 1428-1440.
- Chaban, B., Ng, S. Y., and Jarrell, K. F. (2006). Archaeal habitats--from the extreme to the ordinary. *Can J Microbiol* 52, 73-116.
- Chaudhary, P. P., Brablcová, L., Buriánková, I., and Rulík, M. (2013). Molecular diversity and tools for deciphering the methanogen community structure and diversity in freshwater sediments. *Applied microbiology and biotechnology* 97, 7553-7562.
- Caetano-Anollés, G., Nasir, A., Zhou, K., Caetano-Anollés, D., Mitterthaler, J. E., Sun, F.-J., and Kim, K. M. (2014). Archaea: the first domain of diversified life. *Archaea* 2014.
- Chen, Y., Dumont, M. G., Cébron, A., and Murrell, J. C. (2007). Identification of active methanotrophs in a landfill cover soil through detection of expression of 16S rRNA and functional genes. *Environmental Microbiology* 9, 2855-2869.
- Cheng, Y., Mao, S., Liu, J., and Zhu, W. (2009). Molecular diversity analysis of rumen methanogenic Archaea from goat in eastern China by DGGE methods using different primer pairs. *Letters in applied microbiology* 48, 585-592.
- Chistoserdova, L., Lapidus, A., Han, C., Goodwin, L., Saunders, L., Brettin, T., Tapia, R., Gilna, P., Lucas, S., and Richardson, P. M. (2007). Genome of *Methylobacillus flagellatus*, molecular basis for obligate methylotrophy, and polyphyletic origin of methylotrophy. *Journal of bacteriology* 189, 4020-4027.
- Cho, J.-C., and Tiedje, J. M. (2002). Quantitative detection of microbial genes by using DNA microarrays. *Applied and environmental microbiology* 68, 1425-1430.
- Christensen, T. H., Cossu, R., and Stegmann, R. (1992). Landfilling of waste: leachate, Vol 1: Taylor & Francis).

- Christenson, S., and Cozzarelli, I. (1999). Geochemical and microbiological processes in ground water and surface water affected by municipal landfill leachate. Paper presented at: US Geological Survey Toxic Substances Hydrology Program, in: Proc. of Technical Meeting, Charleston, SC March.
- Conrad, R. (2007). Microbial ecology of methanogens and methanotrophs. *Advances in agronomy* 96, 1-63.
- Cozzarelli, I. M., Suflita, J. M., Ulrich, G. A., Harris, S. H., Scholl, M. A., Schlottmann, J. L., and Christenson, S. (2000). Geochemical and microbiological methods for evaluating anaerobic processes in an aquifer contaminated by landfill leachate. *Environmental science & technology* 34, 4025-4033.
- Cram, D., Sherf, B., Libby, R., Mattaliano, R., Ramachandran, K., and Reeve, J. (1987). Structure and expression of the genes, *mcrBDCGA*, which encode the subunits of component C of methyl coenzyme M reductase in *Methanococcus vannielii*. *Proceedings of the National Academy of Sciences* 84, 3992-3996.
- Das, S., and Dash, H. R. (2013). Molecular Phylogenetics of Microbes. In *Analyzing Microbes*, (Springer), pp. 245-260.
- De Bok, F., Plugge, C., and Stams, A. (2004). Interspecies electron transfer in methanogenic propionate degrading consortia. *Water Research* 38, 1368-1375.
- Desvaux, M., Guedon, E., and Petitdemange, H. (2001). Kinetics and metabolism of cellulose degradation at high substrate concentrations in steady-state continuous cultures of *Clostridium cellulolyticum* on a chemically defined medium. *Applied and environmental microbiology* 67, 3837-3845.
- Dridi, B., Fardeau, M.-L., Ollivier, B., Raoult, D., and Drancourt, M. (2012). *Methanomassiliicoccus luminyensis* gen. nov., sp. nov., a methanogenic archaeon isolated from human faeces. *International journal of systematic and evolutionary microbiology* 62, 1902-1907.

- Dunbar, J., Ticknor, L. O., and Kuske, C. R. (2000). Assessment of microbial diversity in four southwestern United States soils by 16S rRNA gene terminal restriction fragment analysis. *Applied and environmental microbiology* 66, 2943-2950.
- Eichler, J. (2001). Biotechnological uses of archaeal extremozymes. *Biotechnology Advances* 19, 261-278.
- Ellis Joshua, T., Cody, T., and Sims Ronald, C. (2012). Characterization of a methanogenic community within an algal fed anaerobic digester. *ISRN microbiology* 2012.
- Feng, Y., Lin, X., Yu, Y., Zhang, H., Chu, H., and Zhu, J. (2013). Elevated ground-level O₃ negatively influences paddy methanogenic archaeal community. *Scientific reports* 3.
- Galand, P., Fritze, H., Conrad, R., and Yrjälä, K. (2005). Pathways for methanogenesis and diversity of methanogenic archaea in three boreal peatland ecosystems. *Applied and environmental microbiology* 71, 2195-2198.
- García-Maldonado, J. Q., Bebout, B. M., Celis, L. B., and López-Cortés, A. (2012). Phylogenetic diversity of methyl-coenzyme M reductase (mcrA) gene and methanogenesis from trimethylamine in hypersaline environments. *International Microbiology* 15, 33-41.
- Garcia, J.-L., Patel, B. K., and Ollivier, B. (2000). Taxonomic, Phylogenetic, and Ecological Diversity of Methanogenic Archaea. *Anaerobe* 6, 205-226.
- Garrity, G. M., Bell, J. A., and Lilburn, T. G. (2004). Taxonomic outline of the prokaryotes. *Bergey's manual of systematic bacteriology*. Springer, New York, Berlin, Heidelberg.
- Garrity, G. M., and Holt, J. G. (2001). The road map to the manual. In *Bergey's Manual® of Systematic Bacteriology*, (Springer), pp. 119-166.

- Galand, P., Fritze, H., Conrad, R., and Yrjälä, K. (2005). Pathways for methanogenesis and diversity of methanogenic archaea in three boreal peatland ecosystems. *Applied and environmental microbiology* 71, 2195-2198.
- Gerardi, M.H. 2003. Anaerobic digestion stages. *The Microbiology of Anaerobic Digesters*:51-57.
- Godon, J.-J., Zumstein, E., Dabert, P., Habouzit, F., and Moletta, R. (1997). Molecular microbial diversity of an anaerobic digester as determined by small-subunit rDNA sequence analysis. *Appl Environ Microbiol* 63, 2802-2813.
- Gogleva, A. A., Kaparullina, E. N., Doronina, N. V., and Trotsenko, Y. A. (2011). *Methylobacillus arboreus* sp. nov., and *Methylobacillus gramineus* sp. nov., novel non-pigmented obligately methylotrophic bacteria associated with plants. *Systematic and applied microbiology* 34, 477-481.
- Greene, E., Hubert, C., Nemati, M., Jenneman, G., and Voordouw, G. (2003). Nitrite reductase activity of sulphate-reducing bacteria prevents their inhibition by nitrate-reducing, sulphide-oxidizing bacteria. *Environmental Microbiology* 5, 607-617.
- Griggs, D. J., and Noguer, M. (2002). Climate change 2001: the scientific basis. Contribution of working group I to the third assessment report of the intergovernmental panel on climate change. *Weather* 57, 267-269.
- Großkopf, R., Janssen, P. H., and Liesack, W. (1998). Diversity and structure of the methanogenic community in anoxic rice paddy soil microcosms as examined by cultivation and direct 16S rRNA gene sequence retrieval. *Applied and environmental microbiology* 64, 960-969.
- Gupta, S., Choudhary, N., and Alappat, B. J. (2007). Bioreactor landfill for MSW Disposal in Delhi. Paper presented at: Proceedings of the International Conference on Sustainable Solid Waste Management, Chennai, India.

- Ghosh, P., Gupta, A., and Thakur, I. S. (2015). Combined chemical and toxicological evaluation of leachate from municipal solid waste landfill sites of Delhi, India. *Environmental Science and Pollution Research*, 1-11.
- Han, B., Chen, Y., Abell, G., Jiang, H., Bodrossy, L., Zhao, J., Murrell, J. C., and Xing, X. H. (2009). Diversity and activity of methanotrophs in alkaline soil from a Chinese coal mine. *FEMS microbiology ecology* 70, 196-207.
- Handelsman, J. (2004). Metagenomics: application of genomics to uncultured microorganisms. *Microbiology and molecular biology reviews* 68, 669-685.
- Handelsman, J. (2007). Metagenomics and microbial communities. eLS.
- Handelsman, J., Rondon, M. R., Brady, S. F., Clardy, J., and Goodman, R. M. (1998). Molecular biological access to the chemistry of unknown soil microbes: a new frontier for natural products. *Chemistry & Biology* 5, R245-R249.
- Handelsman, J., Tiedje, J., Alvarez-Cohen, L., Ashburner, M., Cann, I., Delong, E., Doolittle, W., Fraser-Liggett, C., Godzik, A., and Gordon, J. (2007). The new science of metagenomics: Revealing the secrets of our microbial planet. *Nat Res Council Report* 13.
- He, Y., Li, Z., Yao, L., Zhao, Y. C., Huang, M. S., and Zhou, G. M. (2014). Molecular phylogenetic analysis of dominant microbial populations in aged refuse. *World Journal of Microbiology and Biotechnology* 30, 1037-1045.
- Hedlund, B. P., Dodsworth, J. A., Cole, J. K., and Panosyan, H. H. (2013). An integrated study reveals diverse methanogens, Thaumarchaeota, and yet-uncultivated archaeal lineages in Armenian hot springs. *Antonie van Leeuwenhoek* 104, 71-82.
- Høj, L., Olsen, R. A., and Torsvik, V. L. (2005). Archaeal communities in High Arctic wetlands at Spitsbergen, Norway (78 N) as characterized by 16S rRNA gene fingerprinting. *FEMS microbiology ecology* 53, 89-101.

- Høj, L., Olsen, R. A., and Torsvik, V. L. (2007). Effects of temperature on the diversity and community structure of known methanogenic groups and other archaea in high Arctic peat. *The ISME journal* 2, 37-48.
- Høj, L., Olsen, R. A., and Torsvik, V. L. (2008). Effects of temperature on the diversity and community structure of known methanogenic groups and other archaea in high Arctic peat. *The ISME journal* 2, 37-48.
- Houghton, J. T., Ding, Y., Griggs, D. J., Noguer, M., van der LINDEN, P. J., Dai, X., Maskell, K., and Johnson, C. (2001). *Climate change 2001: the scientific basis*, Vol 881: Cambridge university press Cambridge).
- Huang, L. N., Chen, Y. Q., Zhou, H., Luo, S., Lan, C. Y., and Qu, L. H. (2003). Characterization of methanogenic Archaea in the leachate of a closed municipal solid waste landfill. *FEMS microbiology ecology* 46, 171-177.
- Huang, L. N., Zhou, H., Zhu, S., and Qu, L. H. (2004). Phylogenetic diversity of bacteria in the leachate of a full-scale recirculating landfill. *FEMS microbiology ecology* 50, 175-183.
- Huang, Y., Zong, W., Yan, X., Wang, R., Hemme, C. L., Zhou, J., and Zhou, Z. (2010). Succession of the bacterial community and dynamics of hydrogen producers in a hydrogen-producing bioreactor. *Applied and environmental microbiology* 76, 3387-3390.
- Huber, H., Hohn, M. J., Stetter, K. O., and Rachel, R. (2003). The phylum Nanoarchaeota: present knowledge and future perspectives of a unique form of life. *Research in microbiology* 154, 165-171.
- Hugenholtz, P., and Tyson, G. W. (2008). Microbiology: metagenomics. *Nature* 455, 481-483.
- Hung, C.-H., Chang, Y.-T., and Chang, Y.-J. (2011). Roles of microorganisms other than *Clostridium* and *Enterobacter* in anaerobic fermentative biohydrogen production systems—a review. *Bioresource technology* 102, 8437-8444.

- Hungate, R. (1969). Chapter IV A Roll Tube Method for Cultivation of Strict Anaerobes. *Methods in microbiology* 3, 117-132.
- Iino, T., Mori, K., and Suzuki, K.-i. (2010). *Methanospirillum lacunae* sp. nov., a methane-producing archaeon isolated from a puddly soil, and emended descriptions of the genus *Methanospirillum* and *Methanospirillum hungatei*. *International journal of systematic and evolutionary microbiology* 60, 2563-2566.
- Ikenaga, M., Asakawa, S., Muraoka, Y., and Kimura, M. (2004). Methanogenic archaeal communities in rice roots grown in flooded soil pots: estimation by PCR-DGGE and sequence analyses. *Soil Science and Plant Nutrition* 50, 701-711.
- Imachi, H., Aoi, K., Tasumi, E., Saito, Y., Yamanaka, Y., Saito, Y., Yamaguchi, T., Tomaru, H., Takeuchi, R., and Morono, Y. (2011). Cultivation of methanogenic community from subseafloor sediments using a continuous-flow bioreactor. *The ISME journal* 5, 1913-1925.
- Iverson, V., Morris, R. M., Frazar, C. D., Berthiaume, C. T., Morales, R. L., and Armbrust, E. V. (2012). Untangling genomes from metagenomes: revealing an uncultured class of marine Euryarchaeota. *Science* 335, 587-590.
- Jain, S., Caforio, A., and Driessen, A. J. (2014). Biosynthesis of archaeal membrane ether lipids. *Frontiers in Microbiology* 5.
- Jarrell, K. F., Walters, A. D., Bochiwal, C., Borgia, J. M., Dickinson, T., and Chong, J. P. (2011). Major players on the microbial stage: why archaea are important. *Microbiology* 157, 919-936.
- Jeyanathan, J., Kirs, M., Ronimus, R. S., Hoskin, S. O., and Janssen, P. H. (2011). Methanogen community structure in the rumens of farmed sheep, cattle and red deer fed different diets. *FEMS microbiology ecology* 76, 311-326.

- Jhamnani, B., and Singh, S. (2009). Groundwater contamination due to Bhalaswa landfill site in New Delhi. *International Journal of Environmental Science and Engineering 1*, 121-125.
- Jiang, H., Chen, Y., Jiang, P., Zhang, C., Smith, T. J., Murrell, J. C., and Xing, X.-H. (2010). Methanotrophs: multifunctional bacteria with promising applications in environmental bioengineering. *Biochemical Engineering Journal 49*, 277-288.
- Juottonen, H. (2008). Archaea, Bacteria, and methane production along environmental gradients in fens and bogs.
- Juottonen, H., Galand, P. E., and Yrjälä, K. (2006). Detection of methanogenic Archaea in peat: comparison of PCR primers targeting the *mcrA* gene. *Research in microbiology 157*, 914-921.
- Juottonen, H., Tuittila, E.-S., Juutinen, S., Fritze, H., and Yrjälä, K. (2008). Seasonality of rDNA-and rRNA-derived archaeal communities and methanogenic potential in a boreal mire. *The ISME journal 2*, 1157-1168.
- Kalia, V. (2008). Microbial treatment of domestic and industrial wastes for bioenergy production.
- Kalyuzhnaya, M. G., Yang, S., Rozova, O. N., Smalley, N. E., Clubb, J., Lamb, A., Gowda, G. A. N., Raftery, D., Fu, Y., Bringel, F., *et al.* (2013). Highly efficient methane biocatalysis revealed in a methanotrophic bacterium. *Nat Commun 4*.
- Khelaifia, S., Raoult, D., and Drancourt, M. (2013). A versatile medium for cultivating methanogenic archaea. *PloS one 8*, e61563.
- Kimura, M. (1980). A simple method for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences. *Journal of molecular evolution 16*, 111-120.

- Kirk, J. L., Beaudette, L. A., Hart, M., Moutoglis, P., Klironomos, J. N., Lee, H., and Trevors, J. T. (2004). Methods of studying soil microbial diversity. *Journal of Microbiological Methods* 58, 169-188.
- Knief, C., Lipski, A., and Dunfield, P. F. (2003). Diversity and activity of methanotrophic bacteria in different upland soils. *Applied and environmental microbiology* 69, 6703-6714.
- Knight, R., Jansson, J., Field, D., Fierer, N., Desai, N., Fuhrman, J. A., Hugenholtz, P., van der Lelie, D., Meyer, F., and Stevens, R. (2012). Unlocking the potential of metagenomics through replicated experimental design. *Nature biotechnology* 30, 513-520.
- Koizumi, Y., Takii, S., and Fukui, M. (2004). Depth-related change in archaeal community structure in a freshwater lake sediment as determined with denaturing gradient gel electrophoresis of amplified 16S rRNA genes and reversely transcribed rRNA fragments. *FEMS microbiology ecology* 48, 285-292.
- Kumar, S., Bhattacharyya, J., Vaidya, A., Chakrabarti, T., Devotta, S., and Akolkar, A. (2009). Assessment of the status of municipal solid waste management in metro cities, state capitals, class I cities, and class II towns in India: an insight. *Waste management* 29, 883-895.
- Kovalic, D., Kwak, J.-H., and Weisblum, B. (1991). General method for direct cloning of DNA fragments generated by the polymerase chain reaction. *Nucleic acids research* 19, 4560-4560.
- Kwok, P.-Y., and Chen, X. (2003). Detection of single nucleotide polymorphisms. *Current issues in molecular biology* 5, 43-60.
- Laloui-Carpentier, W., Li, T., Vigneron, V., Mazeas, L., and Bouchez, T. (2006). Methanogenic diversity and activity in municipal solid waste landfill leachates. *Antonie van Leeuwenhoek* 89, 423-434.

- Lapidus, A., Clum, A., LaButti, K., Kaluzhnaya, M. G., Lim, S., Beck, D. A., del Rio, T. G., Nolan, M., Mavromatis, K., and Huntemann, M. (2011). Genomes of three methylophils from a single niche reveal the genetic and metabolic divergence of the Methylophilaceae. *Journal of bacteriology* *193*, 3757-3764.
- Langille, M. G., Zaneveld, J., Caporaso, J. G., McDonald, D., Knights, D., Reyes, J. A., Clemente, J. C., Burkepille, D. E., Thurber, R. L. V., and Knight, R. (2013). Predictive functional profiling of microbial communities using 16S rRNA marker gene sequences. *Nature biotechnology* *31*, 814-821.
- Lee, C., Kim, J., Shin, S. G., and Hwang, S. (2008). Monitoring bacterial and archaeal community shifts in a mesophilic anaerobic batch reactor treating a high-strength organic wastewater. *FEMS microbiology ecology* *65*, 544-554.
- Lee, C., Kim, J., Shin, S. G., O'Flaherty, V., and Hwang, S. (2010). Quantitative and qualitative transitions of methanogen community structure during the batch anaerobic digestion of cheese-processing wastewater. *Applied microbiology and biotechnology* *87*, 1963-1973.
- Lelieveld, J., Crutzen, P. J., and Dentener, F. J. (1998). Changing concentration, lifetime and climate forcing of atmospheric methane. *Tellus B* *50*, 128-150.
- Liang, C., Xue, Y., Fioroni, M., Rodríguez-Ropero, F., Zhou, C., Schwaneberg, U., and Ma, Y. (2011). Cloning and characterization of a thermostable and halo-tolerant endoglucanase from *Thermoanaerobacter tengcongensis* MB4. *Applied microbiology and biotechnology* *89*, 315-326.
- Liu, C., Zhu, Z., Liu, Y., Guo, T., and Dong, H. (2012a). Diversity and abundance of the rumen and fecal methanogens in Altay sheep native to Xinjiang and the influence of diversity on methane emissions. *Archives of microbiology* *194*, 353-361.
- Liu, D., Ding, W., Jia, Z., and Cai, Z. (2011). Relation between methanogenic archaea and methane production potential in selected natural wetland ecosystems across China. *Biogeosciences* *8*, 329-338.

- Liu, D., Ding, W., Jia, Z., and Cai, Z. (2012b). The impact of dissolved organic carbon on the spatial variability of methanogenic archaea communities in natural wetland ecosystems across China. *Applied microbiology and biotechnology* *96*, 253-263.
- Liu, W.-T., Marsh, T. L., Cheng, H., and Forney, L. J. (1997). Characterization of microbial diversity by determining terminal restriction fragment length polymorphisms of genes encoding 16S rRNA. *Applied and environmental microbiology* *63*, 4516-4522.
- Liu, Y., and Whitman, W. B. (2008). Metabolic, phylogenetic, and ecological diversity of the methanogenic archaea. *Annals of the New York Academy of Sciences* *1125*, 171-189.
- Lloyd, K. G., Schreiber, L., Petersen, D. G., Kjeldsen, K. U., Lever, M. A., Steen, A. D., Stepanauskas, R., Richter, M., Kleindienst, S., and Lenk, S. (2013). Predominant archaea in marine sediments degrade detrital proteins. *Nature* *496*, 215-218.
- Lueders, T., Chin, K. J., Conrad, R., and Friedrich, M. (2001). Molecular analyses of methyl-coenzyme M reductase alpha-subunit (*mcrA*) genes in rice field soil and enrichment cultures reveal the methanogenic phenotype of a novel archaeal lineage. *Environ Microbiol* *3*, 194-204.
- Luton, P. E., Wayne, J. M., Sharp, R. J., and Riley, P. W. (2002). The *mcrA* gene as an alternative to 16S rRNA in the phylogenetic analysis of methanogen populations in landfill. *Microbiology* *148*, 3521-3530.
- Lynch, M. D. J., and Neufeld, J. D. (2015). Ecology and exploration of the rare biosphere. *Nat Rev Micro* *13*, 217-229.
- M., C. S. a. C. I. (1999). Geochemical and microbiological processes in ground water and surface water affected by municipal landfill leachate. . Paper presented at: U.S. Geological Survey Toxic Substances Hydrology Program, Proceedings of the Technical Meeting (Charleston, SC).

- Ma, K., Qiu, Q., and Lu, Y. (2010). Microbial mechanism for rice variety control on methane emission from rice field soil. *Global change biology* *16*, 3085-3095.
- Maarit Niemi, R., Heiskanen, I., Wallenius, K., and Lindström, K. (2001). Extraction and purification of DNA in rhizosphere soil samples for PCR-DGGE analysis of bacterial consortia. *Journal of Microbiological Methods* *45*, 155-165.
- MacNaughton, S. J., Stephen, J. R., Venosa, A. D., Davis, G. A., Chang, Y.-J., and White, D. C. (1999). Microbial population changes during bioremediation of an experimental oil spill. *Applied and environmental microbiology* *65*, 3566-3574.
- Marchuk, D., Drumm, M., Saulino, A., and Collins, F. S. (1991). Construction of T-vectors, a rapid and general system for direct cloning of unmodified PCR products. *Nucleic acids research* *19*, 1154-1154.
- Matte-Tailliez, O., Brochier, C., Forterre, P., and Philippe, H. (2002). Archaeal phylogeny based on ribosomal proteins. *Molecular Biology and Evolution* *19*, 631-639.
- Matthews, G. V. T. (1993). *The Ramsar Convention on Wetlands: its history and development.* (Ramsar convention bureau Gland).
- McDonald, I. R., Upton, M., Hall, G., Pickup, R. W., Edwards, C., Saunders, J. R., Ritchie, D. A., and Murrell, J. C. (1999). Molecular Ecological Analysis of Methanogens and Methanotrophs in Blanket Bog Peat. *Microb Ecol* *38*, 225-233.
- Metje, M., and Frenzel, P. (2005). Effect of temperature on anaerobic ethanol oxidation and methanogenesis in acidic peat from a northern wetland. *Applied and environmental microbiology* *71*, 8191-8200.
- Meuer, J., Kuettner, H. C., Zhang, J. K., Hedderich, R., and Metcalf, W. W. (2002). Genetic analysis of the archaeon *Methanosarcina barkeri* Fusaro reveals a

central role for Ech hydrogenase and ferredoxin in methanogenesis and carbon fixation. *Proceedings of the National Academy of Sciences* 99, 5632-5637.

Milferstedt, K., Youngblut, N. D., and Whitaker, R. J. (2010). Spatial structure and persistence of methanogen populations in humic bog lakes. *The ISME journal* 4, 764-776.

Mitigation, C. C. (2011). IPCC special report on renewable energy sources and climate change mitigation.

Mondav, R., Woodcroft, B. J., Kim, E.-H., McCalley, C. K., Hodgkins, S. B., Crill, P. M., Chanton, J., Hurst, G. B., VerBerkmoes, N. C., and Saleska, S. R. (2014). Discovery of a novel methanogen prevalent in thawing permafrost. *Nature communications* 5.

Morris, R., Schauer-Gimenez, A., Bhattad, U., Kearney, C., Struble, C. A., Zitomer, D., and Maki, J. S. (2014). Methyl coenzyme M reductase (*mcrA*) gene abundance correlates with activity measurements of methanogenic H₂/CO₂-enriched anaerobic biomass. *Microbial biotechnology* 7, 77-84.

Murrell, J. C., McDonald, I. R., and Bourne, D. G. (1998). Molecular methods for the study of methanotroph ecology. *FEMS microbiology ecology* 27, 103-114.

Muyzer, G. (1999). DGGE/TGGE a method for identifying genes from natural ecosystems. *Current opinion in microbiology* 2, 317-322.

Muyzer, G., De Waal, E. C., and Uitterlinden, A. G. (1993). Profiling of complex microbial populations by denaturing gradient gel electrophoresis analysis of polymerase chain reaction-amplified genes coding for 16S rRNA. *Appl Environ Microbiol* 59, 695-700.

Narihiro, T., and Sekiguchi, Y. (2007). Microbial communities in anaerobic digestion processes for waste and wastewater treatment: a microbiological update. *Current opinion in biotechnology* 18, 273-278.

- Narihiro, T., and Sekiguchi, Y. (2011). Oligonucleotide primers, probes and molecular methods for the environmental monitoring of methanogenic archaea. *Microbial biotechnology* 4, 585-602.
- Nemati, M., Mazutinec, T., Jenneman, G., and Voordouw, G. (2001). Control of biogenic H₂S production with nitrite and molybdate. *Journal of Industrial Microbiology and Biotechnology* 26, 350-355.
- Nesme, J., Cécillon, S., Delmont, T. O., Monier, J.-M., Vogel, T. M., and Simonet, P. (2014). Large-Scale Metagenomic-Based Study of Antibiotic Resistance in the Environment. *Current Biology* 24, 1096-1100.
- Nocker, A., Burr, M., and Camper, A. K. (2007). Genotypic microbial community profiling: a critical technical review. *Microbial ecology* 54, 276-289.
- Olsen, G. J., and Woese, C. R. (1993). Ribosomal RNA: a key to phylogeny. *The FASEB journal* 7, 113-123.
- Osborn, A. M., Moore, E. R., and Timmis, K. N. (2000). An evaluation of terminal-restriction fragment length polymorphism (T-RFLP) analysis for the study of microbial community structure and dynamics. *Environmental Microbiology* 2, 39-50.
- Øvreås, L. (2000). Population and community level approaches for analysing microbial diversity in natural environments. *Ecology Letters* 3, 236-251.
- Olsen, G. J., Lane, D. J., Giovannoni, S. J., Pace, N. R., and Stahl, D. A. (1986). Microbial ecology and evolution: a ribosomal RNA approach. *Annual Reviews in Microbiology* 40, 337-365.
- Pace, N., Stahl, D., Lane, D., and Olsen, G. (1986). The analysis of natural microbial populations by ribosomal RNA sequences. *Advances in microbial ecology* 9, 1-55.

- Pace, N. R. (1996). New perspective on the natural microbial world: molecular microbial ecology. *ASM news* 62, 463-470.
- Pace, N. R. (1997). A molecular view of microbial diversity and the biosphere. *Science* 276, 734-740.
- Panigrahy, S., Murthy, T., Patel, J., and Singh, T. (2012). Wetlands of India: inventory and assessment at 1: 50, 000 scale using geospatial techniques. *Current Science* 102, 852-856.
- Parte, A. C. (2013). LPSN—list of prokaryotic names with standing in nomenclature. *Nucleic acids research*, gkt1111.
- Patel, G. B., and Sprott, G. D. (1999). Archaeobacterial ether lipid liposomes (archaeosomes) as novel vaccine and drug delivery systems. *Critical reviews in biotechnology* 19, 317-357.
- Piterina, A. V., and Pembroke, J. T. (2013). Use of PCR-DGGE Based Molecular Methods to Analyse Microbial Community Diversity and Stability during the Thermophilic Stages of an ATAD Wastewater Sludge Treatment Process as an Aid to Performance Monitoring. *ISRN Biotechnology* 2013.
- Prescott, L. M., Harley, J. P., and Klein, D. A. (2002). *Microbiology*. Wm C Brown Publishers.
- Prosser, J. I. (2002). Molecular and functional diversity in soil micro-organisms. *Plant and Soil* 244, 9-17.
- Purvaja, R., and Ramesh, R. (2001). Natural and anthropogenic methane emission from coastal wetlands of South India. *Environmental Management* 27, 547-557.
- Purwantini, E., Torto-Alalibo, T., Lomax, J., Setubal, J. C., Tyler, B. M., and Mukhopadhyay, B. (2014). Genetic resources for methane production from biomass described with gene ontology. *Frontiers in Microbiology* 5.

- Ranjan, M. R., Ramanathan, A., Tripathi, A., and Jha, P. K. (2014). Landfill mining: a case study from Ghazipur landfill area of Delhi. *International Journal of Environmental Sciences* 4, 919-925.
- Rastogi, G., Barua, S., Sani, R. K., and Peyton, B. M. (2011). Investigation of microbial populations in the extremely metal-contaminated Coeur d'Alene River sediments. *Microbial ecology* 62, 1-13.
- Rawat, M., and Ramanathan, A. (2011). Assessment of methane flux from municipal solid waste (MSW) landfill areas of Delhi, India. *Journal of Environmental Protection* 2, 399.
- Reeve, J. N., Nölling, J., Morgan, R. M., and Smith, D. R. (1997). Methanogenesis: genes, genomes, and who's on first? *Journal of bacteriology* 179, 5975.
- Reitschuler, C., Lins, P., and Illmer, P. (2014). Primer evaluation and adaption for cost-efficient SYBR Green-based qPCR and its applicability for specific quantification of methanogens. *World Journal of Microbiology and Biotechnology* 30, 293-304.
- Riesenfeld, C. S., Schloss, P. D., and Handelsman, J. (2004). Metagenomics: genomic analysis of microbial communities. *Annu Rev Genet* 38, 525-552.
- Riviere, D., Desvignes, V., Pelletier, E., Chaussonnerie, S., Guermazi, S., Weissenbach, J., Li, T., Camacho, P., and Sghir, A. (2009). Towards the definition of a core of microorganisms involved in anaerobic digestion of sludge. *The ISME journal* 3, 700-714.
- Rodríguez-Valera, F. (2004). Environmental genomics, the big picture? *FEMS microbiology letters* 231, 153-158.
- Rondon, M. R., August, P. R., Bettermann, A. D., Brady, S. F., Grossman, T. H., Liles, M. R., Loiacono, K. A., Lynch, B. A., MacNeil, I. A., and Minor, C. (2000). Cloning the soil metagenome: a strategy for accessing the genetic and

functional diversity of uncultured microorganisms. *Appl Environ Microbiol* 66, 2541-2547.

Rondon, M. R., Goodman, R. M., and Handelsman, J. (1999). The Earth's bounty: assessing and accessing soil microbial diversity. *Trends in biotechnology* 17, 403-409.

Rother, M., and Metcalf, W. W. (2005). Genetic technologies for Archaea. *Current opinion in microbiology* 8, 745-751.

Roy, S., and Gupta, A. (2012). Water quality assessment of river barak and tributaries in Assam, India. *Pollution Research* 31, 445-450.

Sathaye, J., Shukla, P., and Ravindranath, N. (2006). Climate change, sustainable development and India: Global and national concerns. *CURRENT SCIENCE-BANGALORE*- 90, 314.

Sawayama, S., Tsukahara, K., and Yagishita, T. (2006). Phylogenetic description of immobilized methanogenic community using real-time PCR in a fixed-bed anaerobic digester. *Bioresource technology* 97, 69-76.

Schimel, J. (2000). Global change: rice, microbes and methane. *Nature* 403, 375-377.

Schippers, A., and Neretin, L. N. (2006). Quantification of microbial communities in near-surface and deeply buried marine sediments on the Peru continental margin using real-time PCR. *Environmental Microbiology* 8, 1251-1260.

Schleper, C., Jurgens, G., and Jonscheit, M. (2005). Genomic studies of uncultivated archaea. *Nature Reviews Microbiology* 3, 479-488.

Serrano-Silva, N., Sarria-Guzmán, Y., Dendooven, L., and Luna-Guido, M. (2014). Methanogenesis and Methanotrophy in Soil: A Review. *Pedosphere* 24, 291-307.

Sharma, S., Bhattacharya, S., and Garg, A. (2006). Greenhouse gas emissions from India: a perspective. *Current Science* 90, 326.

- Shin, S. G., Lee, C., Hwang, K., Ahn, J.-H., and Hwang, S. (2008). Use of order-specific primers to investigate the methanogenic diversity in acetate enrichment system. *Journal of industrial microbiology & biotechnology* 35, 1345-1352.
- Singh, J. S. (2011). Methanotrophs: the potential biological sink to mitigate the global methane load. *Current Science* 100, 29-30.
- Singh, V., and Mittal, A. (2011). Toxicity Analysis and Public Health Aspects of Municipal Landfill Leachate: A Case Study of Okhla Landfill, Delhi. Paper presented at: 8th World Wide Workshop for Young Environmental Scientists WWW-YES 2009: Urban waters: resource or risks? 2-5 June 2009.
- Singhal, S., and Pandey, S. (2001). Solid waste management in India: status and future directions. *TERI Information Monitor on Environmental Science* 6, 1-4.
- Singleton, D. R., Furlong, M. A., Rathbun, S. L., and Whitman, W. B. (2001). Quantitative comparisons of 16S rRNA gene sequence libraries from environmental samples. *Applied and environmental microbiology* 67, 4374-4376.
- Sleytr, U. B., Pum, D., and Sára, M. (1997). Advances in S-layer nanotechnology and biomimetics. *Advances in biophysics* 34, 71-79.
- Solomon, S. (2007). *Climate change 2007-the physical science basis: Working group I contribution to the fourth assessment report of the IPCC, Vol 4: Cambridge University Press*.
- Spang, A., Poehlein, A., Offre, P., Zumbärgel, S., Haider, S., Rychlik, N., Nowka, B., Schmeisser, C., Lebedeva, E. V., and Rattei, T. (2012). The genome of the ammonia-oxidizing *Candidatus Nitrososphaera gargensis*: insights into metabolic versatility and environmental adaptations. *Environmental Microbiology* 14, 3122-3145.

- Steinberg, L. M., and Regan, J. M. (2008a). Phylogenetic comparison of the methanogenic communities from an acidic, oligotrophic fen and an anaerobic digester treating municipal wastewater sludge. *Appl Environ Microbiol* *74*, 6663-6671.
- Supaphol, S., Jenkins, S. N., Intomo, P., Waite, I. S., and O'Donnell, A. G. (2011). Microbial community dynamics in mesophilic anaerobic co-digestion of mixed waste. *Bioresource technology* *102*, 4021-4027.
- Tabatabaei, M., Rahim, R. A., Abdullah, N., Wright, A.-D. G., Shirai, Y., Sakai, K., Sulaiman, A., and Hassan, M. A. (2010). Importance of the methanogenic archaea populations in anaerobic wastewater treatments. *Process Biochemistry* *45*, 1214-1225.
- Takai, K., Nakamura, K., Toki, T., Tsunogai, U., Miyazaki, M., Miyazaki, J., Hirayama, H., Nakagawa, S., Nunoura, T., and Horikoshi, K. (2008). Cell proliferation at 122 C and isotopically heavy CH₄ production by a hyperthermophilic methanogen under high-pressure cultivation. *Proceedings of the National Academy of Sciences* *105*, 10949-10954.
- Tamura, K., Stecher, G., Peterson, D., Filipski, A., and Kumar, S. (2013). MEGA6: Molecular Evolutionary Genetics Analysis Version 6.0. *Molecular Biology and Evolution* *30*, 2725-2729.
- Thauer, R. K., Kaster, A.-K., Seedorf, H., Buckel, W., and Hedderich, R. (2008). Methanogenic archaea: ecologically relevant differences in energy conservation. *Nature Reviews Microbiology* *6*, 579-591.
- Theron, J., and Cloete, T. (2000). Molecular techniques for determining microbial diversity and community structure in natural environments. *Critical Reviews in Microbiology* *26*, 37-57.
- Tiedje, J. M., Asuming-Brempong, S., Nüsslein, K., Marsh, T. L., and Flynn, S. J. (1999). Opening the black box of soil microbial diversity. *Applied Soil Ecology* *13*, 109-122.

- Torsvik, V., and Øvreås, L. (2002). Microbial diversity and function in soil: from genes to ecosystems. *Current opinion in microbiology* 5, 240-245.
- Trends, G., and Framework, S. (2010). Waste AND Climate Change.
- Tringe, S. G., and Rubin, E. M. (2005). Metagenomics: DNA sequencing of environmental samples. *Nature reviews genetics* 6, 805-814.
- V Wintzingerode, F., Göbel, U. B., and Stackebrandt, E. (1997). Determination of microbial diversity in environmental samples: pitfalls of PCR-based rRNA analysis. *FEMS microbiology reviews* 21, 213-229.
- Van Dyke, M., and McCarthy, A. (2002). Molecular biological detection and characterization of *Clostridium* populations in municipal landfill sites. *Applied and environmental microbiology* 68, 2049-2053.
- van Elsas, J. D., Duarte, G. F., Keijzer-Wolters, A., and Smit, E. (2000). Analysis of the dynamics of fungal communities in soil via fungal-specific PCR of soil DNA followed by denaturing gradient gel electrophoresis. *Journal of Microbiological Methods* 43, 133-151.
- Venter, J. C., Remington, K., Heidelberg, J. F., Halpern, A. L., Rusch, D., Eisen, J. A., Wu, D., Paulsen, I., Nelson, K. E., and Nelson, W. (2004). Environmental genome shotgun sequencing of the Sargasso Sea. *Science* 304, 66-74.
- Vigneron, A., Cruaud, P., Pignet, P., Caprais, J.-C., Cambon-Bonavita, M.-A., Godfroy, A., and Toffin, L. (2013). Archaeal and anaerobic methane oxidizer communities in the Sonora Margin cold seeps, Guaymas Basin (Gulf of California). *The ISME journal* 7, 1595-1608.
- Voordouw, G., Shen, Y., Harrington, C. S., Telang, A. J., Jack, T. R., and Westlake, D. W. (1993). Quantitative reverse sample genome probing of microbial communities and its application to oil field production waters. *Applied and environmental microbiology* 59, 4101-4114.

- Watanabe, T., Kimura, M., and Asakawa, S. (2006). Community structure of methanogenic archaea in paddy field soil under double cropping (rice–wheat). *Soil Biology and Biochemistry* 38, 1264-1274.
- Watanabe, T., Kimura, M., and Asakawa, S. (2007). Dynamics of methanogenic archaeal communities based on rRNA analysis and their relation to methanogenic activity in Japanese paddy field soils. *Soil Biology and Biochemistry* 39, 2877-2887.
- Watanabe, T., Kimura, M., and Asakawa, S. (2010). Diversity of methanogenic archaeal communities in Japanese paddy field ecosystem, estimated by denaturing gradient gel electrophoresis. *Biology and fertility of soils* 46, 343-353.
- Whitman, W. B., Bowen, T. L., and Boone, D. R. (2006). The methanogenic bacteria. In *The prokaryotes*, (Springer), pp. 165-207.
- Willey, J. (2008). Prescott, Harley, and Klein's *Microbiology-7th international ed./Joanne M. Willey, Linda M. Sherwood, Christopher J. Woolverton*: New York [etc.]: McGraw-Hill Higher Education).
- Williamson, S. J., and Yooseph, S. (2012). From bacterial to microbial ecosystems (metagenomics). In *Bacterial Molecular Networks*, (Springer), pp. 35-55.
- Woese, C. R., and Fox, G. E. (1977). Phylogenetic structure of the prokaryotic domain: the primary kingdoms. *Proceedings of the National Academy of Sciences* 74, 5088-5090.
- Woese, C. R. (1987). Bacterial evolution. *Microbiol Rev* 51, 221.
- Woese, C. R., Kandler, O., and Wheelis, M. L. (1990). Towards a natural system of organisms: proposal for the domains Archaea, Bacteria, and Eucarya. *Proceedings of the National Academy of Sciences* 87, 4576-4579.

- Wolfe, R. S., Amy, C., and Stephen, W. (2011). Techniques for cultivating methanogens. *Methods in enzymology* *494*, 1-22.
- Yarza, P., Richter, M., Peplies, J., Euzéby, J., Amann, R., Schleifer, K.-H., Ludwig, W., Glöckner, F. O., and Rosselló-Móra, R. (2008). The All-Species Living Tree project: A 16S rRNA-based phylogenetic tree of all sequenced type strains. *Systematic and applied microbiology* *31*, 241-250.
- Yarza, P., Yilmaz, P., Pruesse, E., Glöckner, F. O., Ludwig, W., Schleifer, K.-H., Whitman, W. B., Euzéby, J., Amann, R., and Rosselló-Móra, R. (2014). Uniting the classification of cultured and uncultured bacteria and archaea using 16S rRNA gene sequences. *Nature Reviews Microbiology* *12*, 635-645.
- Yashiro, Y., Sakai, S., Ehara, M., Miyazaki, M., Yamaguchi, T., and Imachi, H. (2011). *Methanoregula formicica* sp. nov., a methane-producing archaeon isolated from methanogenic sludge. *International journal of systematic and evolutionary microbiology* *61*, 53-59.
- Yu, Y., Lee, C., Kim, J., and Hwang, S. (2005). Group-specific primer and probe sets to detect methanogenic communities using quantitative real-time polymerase chain reaction. *Biotechnology and bioengineering* *89*, 670-679.
- Yvon-Durocher, G., Allen, A. P., Bastviken, D., Conrad, R., Gudasz, C., St-Pierre, A., Thanh-Duc, N., and del Giorgio, P. A. (2014). Methane fluxes show consistent temperature dependence across microbial to ecosystem scales. *Nature* *507*, 488-491.
- Zhang, D.-C., Mörtelmaier, C., and Margesin, R. (2012). Characterization of the bacterial archaeal diversity in hydrocarbon-contaminated soil. *Science of the Total Environment* *421*, 184-196.