



Department of Biotechnology

School of Life sciences

Assam University, Silchar

(A Central University Established by an Act of Parliament)

Silchar-788 011, Assam, India

Date:

DECLARATION

I, Arif Uddin, bearing Registration No. **PhD/2423/13**, dated **12/09/13**, hereby declare that the subject matter of the thesis entitled “**Analysis of Codon Usage Bias in Mitochondrial DNA of Some Chordates**”, is the record of work done by me and that the contents of this thesis did not form the basis for an award of any degree to me or to anybody else to the best of my knowledge. The thesis has not been submitted to any other University/Institute.

Place:

(Arif Uddin)

Date:

Research Scholar

ACKNOWLEDGEMENTS

First of all, I thank the almighty Allah the most gracious and the most merciful, for the blessing upon me during my research work, and indeed, throughout my life.

I would like to express my deep sense of gratitude and heart-felt thanks to my supervisor Prof (Dr.) Supriyo Chakraborty for his continuous assistance, guidance, support, suggestions, necessary criticism and encouragement. I honestly appreciate the time and effort he has given me to conduct my research work.

I am also thankful to Dean, Hargobind Khurana School of Life Sciences, Department of Biotechnology and all the faculty members and official staffs of Department of Biotechnology for their kind cooperation as they have always extended their helping hands whenever needed. I would like to express my deepest gratitude to **University Grant Commission (UGC)**, Govt. of India for awarding me Maulana Azad National Fellowship (**MANF**) for pursuing my PhD work and **Department of Biotechnology**, Assam University for providing infrastructural support.

I would like to express my gratitude to my lab mates and friends who had made my university life more enjoyable and pleasant by sharing their invaluable time. Specially I am thankful to Tarikul Huda Mazumder, Monisha Nath Choudhury, Prosenjit Paul, Binata Halder, Himanghu Deka, Arup Malakar and Gulshana A. Mazumder for their support, encouragement and advice.

I am thankful to Fazlur Rahman Talukder, Md. Gulzar Hussain, Rabinson C. Jose, Alanur Hussain Laskar, Ruhina Sirin Laskar, Md. Jaynal Abedin, A.K.Md.Fuzail and Mahfuj Alom Mazumdar for their support, encouragement and advice.

Finally, and most importantly, I would like to acknowledge my beloved parents and family members for their immense mental support, invaluable love and inspiration throughout my academic endeavour.

Dedicated to

Prophet Hazrat Muhammad Mustafa Sallallahu Alaihi wa Sallam,

Parents, Teachers and Hardworking people of our country

CONTENTS

| CHAPTER | PAGE NO. |
|--|-----------------|
| List of Tables | I-III |
| List of Figures | IV-XIII |
| Chapter -1 Introduction | 1-20 |
| 1.1 Genetic code | 1 |
| 1.2 Properties of genetics code | 1-2 |
| 1.3 Codon Usage Bias | 2-12 |
| 1.3.1 Factor for codon usage bias | 3-5 |
| 1.3.1.1 Mutational Bias | 3 |
| 1.3.1.2 Natural selection | 3 |
| 1.3.1.3 Gene expression level | 3-4 |
| 1.3.2 Parameters for codon usage bias | 5-6 |
| 1.3.2.1 Base composition at silent sites | 5 |
| 1.3.2.2 Relative Synonymous Codon Usage (RSCU) | 6 |
| 1.3.2.3 Effective Number of Codons (ENC) | 6 |
| 1.3.2.4 Codon adaptation index (CAI) | 6 |
| 1.3.3 Application of codon usage bias | 6-12 |
| 1.3.3.1 Cotranslational Protein Folding | 7-9 |
| 1.3.3.2 Cotranslational Interactions with Other Cellular Components | 9 |
| 1.3.3.3 Functional Regulation of Expression Level | 9-10 |
| 1.3.3.4 Viral Codon Usage | 10 |
| 1.3.3.5 Codon Usage and Human Health | 10-11 |
| 1.3.3.6 Codon Usage in Horizontally Transferred Genes | 11 |

| | |
|--|--------------|
| 1.3.3.7 Heterologous Protein Expression | 11-12 |
| 1.4 Mitochondrial DNA | 12-16 |
| 1.4.1 Mitochondrial genetic code | 13-14 |
| 1.4.2 Evolution of Mitochondria | 14-15 |
| 1.4.3 Functional Role of Mitochondrial Genomes | 15-16 |
| 1.4.4 Base composition in Mitochondria | 16 |
| 1.5 Chordates | 16-18 |
| 1.5.1 Reason for selection of pisces, aves and Mammals | 17-18 |
| 1.6 Statement of the problem | 19 |
| 1.6.1 Rationale of the study | 19 |
| 1.6.2 Novelty of the Study | 19 |
| 1.7. Objectives | 20 |
| Chapter-2 Review of Literature | 21-33 |
| Chapter-3 Materials and Methodology | 34-37 |
| 3.1 Availability of sequences data | 34 |
| 3.2 Compositional constraints | 34-35 |
| 3.3 Indices of synonymous codon usage bias | 35-36 |
| 3.3.1 Relative synonymous codon usage (RSCU) | 35 |
| 3.3.2 Effective number of codons (ENC) | 35-36 |
| 3.3.3 Codon adaptation index (CAI) | 36 |
| 3.4 Correspondence analysis | 36 |
| 3.5 Neutrality plot | 36-37 |
| 3.6 Hierarchal Clustering | 37 |
| 3.7 Software used | 37 |

Chapter-4 Results **38-149**

| | |
|---|---------------|
| 4.1. Analysis of codon usage bias in MT-ND gene among Pisces, Aves and Mammals | 38-95 |
| 4.1.1 Codon usage pattern in MT-ND gene among pisces, aves and mammals | 38-45 |
| 4.1.2 Analysis of codon usage bias in mitochondrial ND gene | 45-49 |
| 4.1.3 Prediction of expression level for mitochondrial ND gene | 49-52 |
| 4.1.4. Relationships of gene expression (CAI) with codon usage bias (ENC) for MT-ND gene among pisces, aves and mammals | 53-54 |
| 4.1.5 Compositional features of MT-ATP gene among pisces, aves and mammals | 54-66 |
| 4.1.6 Interrelationships among different compositional features of MT-ND gene among pisces, aves and mammals | 66-84 |
| 4.1.7 Correspondence analysis (COA) of MT-ND gene among pisces, aves and mammals | 84-91 |
| 4.1.8 Amino acid composition of ND gene in different species of pisces, aves and mammals | 91-95 |
| 4.2 Analysis of codon usage bias in MT-CO gene among Pisces, Aves and Mammals | 95-118 |
| 4.2.1 Codon usage pattern of CO gene in pisces, aves and mammals | 95-97 |
| 4.2.2 Analysis of codon usage bias in mitochondrial CO | 98-99 |
| 4.2.3 Prediction of expression level for mitochondrial CO gene | 99-101 |
| 4.2.4 Relationships of gene expression (CAI) with codon usage bias for mitochondrial CO gene among Pisces, aves and mammals | 101 |
| 4.2.5 Compositional features of mitochondrial CO gene among pisces, aves and mammals | 102-106 |
| 4.2.6. Interrelationships among different compositional features of MT-CO gene among pisces, aves and mammals | 106-113 |
| 4.2.7. Correspondence analysis (COA) of mitochondrial CO gene among pisces, aves and mammals | 114-116 |
| 4.2.8 Amino acid composition CO gene among pisces, aves and mammals | 117-118 |

| | |
|--|----------------|
| 4.3. Analysis of codon usage bias in MT-CYB gene among Pisces, | |
| Aves and Mammals | 118-127 |
| 4.2.1 Codon usage pattern of CYB gene in pisces, aves and mammals | 118-119 |
| 4.2.2 Analysis of codon usage bias in mitochondrial CYB | 119 |
| 4.2.3 Prediction of expression level for mitochondrial CYB gene | 119-120 |
| 4.2.4 Relationships of gene expression (CAI) with codon usage bias for mitochondrial CYB among Pisces, aves and mammals | 120 |
| 4.2.5 Compositional features of mitochondrial CYB gene among pisces, aves and mammals | 120-122 |
| 4.2.6. Interrelationships among different compositional features of CYB gene among pisces, aves and mammals | 122-125 |
| 4.2.7. Correspondence analysis (COA) of mitochondrial CYB gene among pisces, aves and mammals | 125-126 |
| 4.2.8 Amino acid composition CYB gene among pisces, aves and mammals | 126-127 |
| 4.4 Analysis of codon usage bias in MT-ATP gene among Pisces, | |
| Aves and Mammals | 127-141 |
| 4.2.1 Codon usage pattern of ATP gene in pisces, aves and mammals | 127-128 |
| 4.2.2 Analysis of codon usage bias in mitochondrial ATP | 128-129 |
| 4.2.3 Prediction of expression level for mitochondrial ATP gene | 129-130 |
| 4.2.4 Relationships of gene expression (CAI) with codon usage bias (ENC) for mitochondrial ATP among Pisces, aves and mammals | 130 |
| 4.2.5 Compositional features of mitochondrial ATP gene among pisces, aves and mammals | 131-133 |
| 4.2.6. Interrelationships among different compositional features of ATP gene among pisces, aves and mammals | 134-138 |
| 4.2.7. Correspondence analysis (COA) of mitochondrial ATP gene among pisces, aves and mammals | 138-140 |
| 4.2.8 Amino acid composition ATP gene among pisces, aves and mammals | 141 |
| 4.5 Comparative analysis of codon usage pattern in mitochondrial genes among the chordates | 142-149 |

| | |
|---|-----------------|
| Chapter-5 Discussion | 150-165 |
| 5.1 Codon usage pattern | 151-152 |
| 5.2 Coon usage bias | 153 |
| 5.3 Expression level | 154 |
| 5.4 Interrelationships of gene expression with codon usage bias | 155 |
| 5.5 Compositional features | 155-159 |
| 5.6 Interrelationships among different compositional features | 159-161 |
| 5.7 Correspondence analysis | 161-162 |
| 5.8 Amino acid composition | 162-165 |
| Summary | 166-169 |
| Bibliography | 170-183 |
| Appendix-I List of Abbreviations | XIV-XV |
| Appendix-II List of Publications | XVI-XVII |
| (a) Scientific Publication | XVI |
| (b) Conference Proceedings | XVII |

List of Tables

| Sl No. | Table No. | Title of the table | Page No. |
|--------|--------------|---|----------|
| 1 | Table 3.1. | Accession numbers of coding sequences (cds) of 15 species with their respective family | 34 |
| | Table 4.1.1 | Nucleotide compositions among pisces, aves and mammals for ND1 gene | 55 |
| | Table 4.1.2 | Nucleotide compositions among pisces, aves and mammals for ND2 gene | 57 |
| | Table 4.1.3 | Nucleotide compositions among pisces, aves and mammals for ND3 gene | 59 |
| | Table 4.1.4 | Nucleotide compositions among pisces, aves and mammals for ND4 gene | 61 |
| | Table 4.1.5 | Nucleotide compositions among pisces, aves and mammals for ND4L gene | 62 |
| | Table 4.1.6 | Nucleotide compositions among pisces, aves and mammals for ND5 gene | 64 |
| | Table 4.1.7 | Nucleotide compositions among pisces, aves and mammals for ND6 gene | 66 |
| | Table 4.1.8 | Correlation between overall nucleotide composition (%) and its composition at 3 rd codon position in pisces, aves and mammals in ND1 gene | 67 |
| | Table 4.1.9 | Correlation coefficients among ENC, CAI, GC contents for MT-ND1 gene | 67 |
| | Table 4.1.10 | Correlation between overall nucleotide composition (%) and its composition at 3 rd codon position in pisces, aves and mammals in ND2 gene | 69 |
| | Table 4.1.11 | Correlation coefficients among ENC, CAI, GC contents for MT-ND2 gene | 70 |
| | Table 4.1.12 | Correlation between overall nucleotide composition (%) and its composition at 3 rd codon position in pisces, aves and mammals in ND3 gene | 72 |
| | Table 4.1.13 | Correlation coefficients among ENC, CAI, GC contents for MT-ND3 gene | 72 |
| | Table 4.1.14 | Correlation between overall nucleotide composition (%) and its composition at 3 rd codon position in pisces, aves and mammals in ND4 gene | 74 |
| | Table 4.1.15 | Correlation coefficients among ENC, CAI, GC contents for MT-ND4 gene | 75 |
| | Table 4.1.16 | Correlation between overall nucleotide composition (%) and its composition at 3 rd codon position in pisces, aves and mammals in ND4L gene | 77 |
| | Table 4.1.17 | Correlation coefficients among ENC, CAI, GC contents for MT-ND4L gene | 77 |
| | Table 4.1.18 | Correlation between overall nucleotide composition (%) and its composition at 3 rd codon position in pisces, aves and mammals for ND5 gene | 79 |
| | Table | Correlation coefficients among ENC, CAI, GC contents for MT-ND5 gene | 80 |

| | | | |
|--|--------------|---|-----|
| | 4.1.19 | | |
| | Table 4.1.20 | Correlation between overall nucleotide composition (%) and its composition at 3 rd codon position in pisces, aves and mammals in ND6 gene | 82 |
| | Table 4.1.21 | Correlation coefficients among ENC, CAI, GC contents for MT-ND6 gene | 82 |
| | Table 4.2.1 | Nucleotide compositions among pisces, aves and mammals for COI gene | 103 |
| | Table 4.2.2 | Nucleotide compositions among pisces, aves and mammals for COII gene | 105 |
| | Table 4.2.3 | Nucleotide compositions among pisces, aves and mammals for COIII gene | 106 |
| | Table 4.2.4 | Correlation between overall nucleotide composition (%) and its composition at 3 rd codon position in pisces, aves and mammals for COI gene | 107 |
| | Table 4.2.5 | Correlation coefficients among ENC, CAI, GC contents for COI gene | 107 |
| | Table 4.2.6 | Correlation between overall nucleotide composition (%) and its composition at 3 rd codon position in pisces, aves and mammals for COII gene | 109 |
| | Table 4.2.7 | Correlation coefficients among ENC, CAI, GC contents for COII gene | 110 |
| | Table 4.2.8 | Correlation between overall nucleotide composition (%) and its composition at 3 rd codon position in pisces, aves and mammals for COIII gene | 112 |
| | Table 4.2.9 | Correlation coefficients among ENC, CAI, GC contents for COIII gene | 112 |
| | Table 4.3.1 | Nucleotide compositions among pisces, aves and mammals for CYB gene | 122 |
| | Table 4.3.2 | Correlation between overall nucleotide composition (%) and its composition at 3 rd codon position in pisces, aves and mammals | 123 |
| | Table 4.3.3 | Correlation coefficients among ENC, CAI, GC contents for MT-CYB gene | 123 |
| | Table 4.4.1 | Nucleotide compositions among pisces, aves and mammals for ATP6 gene | 132 |
| | Table 4.4.2 | Nucleotide compositions among pisces, aves and mammals for ATP8 gene | 133 |
| | Table 4.4.3 | Correlation between overall nucleotide composition (%) and its composition at 3 rd codon position in pisces, aves and mammals for ATP6 gene | 134 |
| | Table 4.4.4 | Correlation coefficients among ENC, CAI, GC contents | 135 |
| | Table 4.4.5 | Correlation between overall nucleotide composition (%) and its composition at 3 rd codon position in pisces, aves and Mammals for ATP8 gene | 136 |
| | Table 4.4.6 | Correlation coefficients among ENC, CAI, GC contents for ATP8 gene | 137 |
| | Table | RSCU values of codons for different mitochondrial genes in ND gene | 143 |

| | | | |
|--|-------------|---|---------|
| | 4.5.1 | among pisces, aves and mammals | |
| | Table 4.5.2 | RSCU values of codons for different mitochondrial genes in ND gene among pisces, aves and mammals | 145 |
| | Table 4.5.3 | RSCU values of codons for different mitochondrial genes in CO and CYB gene among pisces, aves and mammals | 147 |
| | Table 4.5.4 | RSCU values of codons for different mitochondrial genes in ATP gene among pisces, aves and mammals | 148-149 |
| | Table 5.1 | ENC values of 13 mitochondrial protein-coding genes among pisces, aves and mammals | 153 |
| | Table 5.2 | CAI values of 13 mitochondrial protein coding genes among pisces, aves and mammals | 154 |
| | Table 5.3 | Comparison of compositional properties in 13 mitochondrial protein-coding genes | 156-158 |
| | Table 5.4 | F1 and F2 of COA among pisces, aves and mammals | 162 |

List of Figures

| Sl No. | Figure No. | Title of the figure | Page No. |
|--------|---------------|--|----------|
| | Figure 1.1 | Rare codons are hypothesized to introduce translational pauses that modulate cotranslational protein folding | 8 |
| | Figure 1.2 | Hypothetical example of heterologous expression of a human coding sequence in <i>Escherichia coli</i> . | 12 |
| | Figure 1.3 | Structure of mitochondrial DNA | 13 |
| | Figure 1.4 | Mitochondrial genetic code | 14 |
| | Figure 1.5 | Different species of chordates under study | 18 |
| | Figure 4.1.1 | Hierarchical Clustering of RSCU values of each codon among pisces, aves and mammals for ND1 gene | 38 |
| | Figure 4.1.2. | Hierarchical Clustering of RSCU values of each codon among pisces, aves and mammals for ND2 gene | 39 |
| | Figure 4.1.3 | Hierarchical Clustering of RSCU values of each codon among pisces, aves and mammals for ND3 gene | 40 |
| | Figure 4.1.4 | Hierarchical Clustering of RSCU values of each codon among pisces, aves and mammals for ND4 gene | 41 |
| | Figure 4.1.5 | Hierarchical Clustering of RSCU values of each codon among pisces, aves and mammals for ND4l gene | 42 |
| | Figure 4.1.6 | Hierarchical Clustering of RSCU values of each codon among pisces, aves and mammals for ND5 gene | 43 |
| | Figure 4.1.7 | Hierarchical Clustering of RSCU values of each codon among pisces, aves and mammals for ND6 gene | 44 |
| | Figure 4.1.8 | Distribution of ENC in different species of pisces, aves and mammals for MT-ND1 gene | 45 |
| | Figure 4.1.9 | Distribution of ENC in different species of pisces, aves and mammals for MT-ND2 gene | 46 |
| | Figure 4.1.10 | Distribution of ENC in different species of pisces, aves and mammals for MT-ND3 gene | 46 |
| | Figure 4.1.11 | Distribution of ENC in different species of pisces, aves and | 47 |

| | | | |
|--|---------------|---|----|
| | | mammals for MT-ND4 gene | |
| | Figure 4.1.12 | Distribution of ENC in different species of pisces, aves and mammals for MT-ND4l gene | 47 |
| | Figure 4.1.13 | Distribution of ENC in different species of pisces, aves and mammals for MT-ND5 gene | 48 |
| | Figure 4.1.14 | Distribution of ENC in different species of pisces, aves and mammals for MT-ND6 gene | 48 |
| | Figure 4.1.15 | Expression level in different species of pisces, aves and mammals for MT-ND1 gene | 49 |
| | Figure 4.1.16 | Expression level in different species of pisces, aves and mammals for MT-ND2 gene | 50 |
| | Figure 4.1.17 | Expression level in different species of pisces, aves and mammals for MT-ND3 gene | 50 |
| | Figure 4.1.18 | Expression level in different species of pisces, aves and mammals for MT-ND4 gene | 51 |
| | Figure 4.1.19 | Expression level in different species of pisces, aves and mammals for MT-ND4l gene | 51 |
| | Figure 4.1.20 | Expression level in different species of pisces, aves and mammals for MT-ND5 gene | 52 |
| | Figure 4.1.21 | Expression level in different species of pisces, aves and mammals for MT-ND6 gene | 52 |
| | Figure 4.1.22 | Nucleotide composition and its composition at 3 rd codon position for MT-ND1 gene | 54 |
| | Figure 4.1.23 | Overall GC content and GC content at 1 st , 2 nd and 3 rd position in pisces, aves and mammals for MT-ND1 gene | 55 |
| | Figure 4.1.24 | Nucleotide composition and its composition at 3 rd codon position for MT-ND2 gene | 56 |
| | Figure 4.1.25 | Overall GC content and GC content at 1 st , 2 nd and 3 rd position in pisces, aves and mammals for MT-ND2 gene | 56 |
| | Figure 4.1.26 | Nucleotide composition and its composition at 3 rd codon position for MT-ND3 gene | 58 |
| | Figure 4.1.27 | Overall GC content and GC content at 1 st , 2 nd and 3 rd position in pisces, aves and mammals for MT-ND3 gene | 58 |

| | | |
|-------------------|--|----|
| Figure 4.1.28 | Overall nucleotide composition and its composition at 3 rd codon position for MT-ND4 gene | 60 |
| Figure 4.1.29 | Overall GC content and GC content at 1 st , 2 nd and 3 rd position in pisces, aves and mammals for MT-ND4 gene | 60 |
| Figure 4.1.30 | Overall nucleotide composition and its composition at 3 rd codon position for MT-ND4l gene | 61 |
| Figure 4.1.31 | Overall GC content and GC content at 1 st , 2 nd and 3 rd position in pisces, aves and mammals for MT-ND4l gene | 62 |
| Figure 4.1.32 | Overall Nucleotide composition and its composition at 3 rd codon position for MT-ND5 gene | 63 |
| Figure 4.1.33 | Overall GC content and GC content at 1 st , 2 nd and 3 rd position in pisces, aves and mammals for MT-ND5 gene | 64 |
| Figure 4.1.34 | Overall nucleotide composition and its composition at 3 rd codon position for MT-ND6 gene | 65 |
| Figure 4.1.35 | GC content and its 1 st , 2 nd and 3 rd position in pisces, aves and mammals for MT-ND6 gene | 65 |
| Figure 4.1.36 (a) | Neutrality plot of GC12 with GC3 in different species of pisces for ND1 gene | 68 |
| Figure 4.1.36 (b) | Neutrality plot of GC12 with GC3 in different species of aves for ND1 gene | 68 |
| Figure 4.1.36 (c) | Neutrality plot of GC12 with GC3 in different species of mammals for ND1 gene | 69 |
| Figure 4.1.37(a) | Neutrality plot of GC12 with GC3 in different species of pisces for ND2 gene | 71 |
| Figure 4.1.37(b) | Neutrality plot of GC12 with GC3 in different species of aves for ND2 gene | 71 |
| Figure 4.1.37 (c) | Neutrality plot of GC12 with GC3 in different species of mammals for ND2 gene | 71 |
| Figure 4.1.38(a) | Neutrality plot of GC12 with GC3 in different species of pisces for ND3 gene | 73 |
| | Fig.4.1.38(b) Neutrality plot of GC12 with GC3 in different species of aves for ND3 gene | |

| | | |
|-------------------|---|----|
| Figure 4.1.38 (c) | Neutrality plot of GC12 with GC3 in different species of mammals for ND3 gene | 73 |
| Figure 4.1.39(a) | Neutrality plot of GC12 with GC3 in different species of pisces for ND4 gene | 74 |
| Figure 4.1.39(b) | Neutrality plot of GC12 with GC3 in different species of aves for ND4 gene | 76 |
| Figure 4.1.39(c) | Neutrality plot of GC12 with GC3 in different species of mammals for ND4 gene | 76 |
| Figure 4.1.40(a) | Neutrality plot of GC12 with GC3 in different species of pisces for ND4L gene | 76 |
| Figure 4.1.40(b) | Neutrality plot of GC12 with GC3 in different species of aves for ND4L gene | 78 |
| Figure 4.1.40 (c) | Neutrality plot of GC12 with GC3 in different species of mammals for ND4L gene | 78 |
| Figure 4.1.41(a) | Neutrality plot of GC12 with GC3 in different species of pisces for ND5 gene | 79 |
| Figure 4.1.41(b) | Neutrality plot of GC12 with GC3 in different species of aves for ND5 gene | 81 |
| Figure 4.1.41(c) | Neutrality plot of GC12 with GC3 in different species of mammals for ND5 gene | 81 |
| Figure 4.1.42(a) | Neutrality plot of GC12 with GC3 in different species of pisces for ND6 gene | 81 |
| Figure 4.1.42(b) | Neutrality plot of GC12 with GC3 in different species of aves for ND6 gene | 83 |
| Figure 4.1.42(c) | Neutrality plot of GC12 with GC3 in different species of mammals for ND6 gene | 83 |
| Figure 4.1.43 | Contributions of the axes for MT-ND1 gene in pisces, aves and mammals | 84 |
| Figure 4.1.44 | Correspondence analysis of the synonymous codon usage towards the codons in ND1 gene for pisces, aves and mammals | 84 |
| Figure 4.1.45 | Contributions of the axes for MT-ND2 gene in pisces, aves and mammals | 85 |

| | | |
|---------------|--|----|
| Figure 4.1.46 | Correspondence analysis of synonymous codon usage towards the codons in ND2 gene for pisces, aves and mammals | 85 |
| Figure 4.1.47 | Contributions of the axes for MT-ND3 gene in pisces, aves and mammals | 86 |
| Figure 4.1.48 | Correspondence analysis of synonymous codon usage towards the codons in ND3 gene for pisces, aves and mammals | 86 |
| Figure 4.1.49 | Contributions of the axes for MT-ND4 gene in pisces, aves and mammals | 87 |
| Figure 4.1.50 | Correspondence analysis of synonymous codon usage towards the codons in ND4 gene for pisces, aves and mammals | 87 |
| Figure 4.1.51 | Contributions of the axes for MT-ND4L gene in pisces, aves and mammals | 88 |
| Figure 4.1.52 | Correspondence analysis of synonymous codon usage towards the codons in ND4L gene for pisces, aves and mammals | 88 |
| Figure 4.1.53 | Contributions of the axes for MT-ND5 gene in pisces, aves and mammals | 89 |
| Figure 4.1.54 | Correspondence analysis of synonymous codon usage towards the codons in ND5 gene for pisces, aves and mammals | 89 |
| Figure 4.1.55 | Contributions of the axes for MT-ND6 gene in pisces, aves and mammals | 90 |
| Figure 4.1.56 | Correspondence analysis of synonymous codon usage towards the codons in ND6 gene for pisces, aves and mammals | 90 |
| Figure 4.1.57 | Amino acid distribution in different species of pisces, aves and mammals for ND1 gene | 91 |
| Figure 4.1.58 | Amino acid distribution in different species of pisces, aves and mammals for ND2 gene | 92 |
| Figure 4.1.59 | Amino acid distribution in different species of pisces, aves and mammals for ND3 gene | 92 |
| Figure 4.1.60 | Amino acid distribution in different species of pisces, aves and mammals for ND4 gene | 93 |
| Figure 4.1.61 | Amino acid distribution in different species of pisces, aves and mammals for ND4L gene | 93 |

| | | |
|-----------------|---|-----|
| Fig. 4.1.62 | Amino acid distribution in different species of pisces, aves and mammals for ND5 gene | 94 |
| Figure 4.1.63 | Amino acid distribution in different species of pisces, aves and mammals for ND6 gene | 94 |
| Figure 4.2.1 | Hierarchical Clustering of RSCU values of each codon among pisces, aves and mammals for COI | 95 |
| Figure 4.2.2 | Hierarchical Clustering of RSCU values of each codon among pisces, aves and mammals for COII gene | 95 |
| Figure 4.2.3 | Hierarchical Clustering of RSCU values of each codon among pisces, aves and mammals for COIII gene | 96 |
| Figure 4.2.4 | Distribution of ENC in different species in pisces, aves and mammals for MT-COI gene | 97 |
| Figure 4.2.5 | Distribution of ENC in different species in pisces, aves and mammals for MT-COII gene | 98 |
| Figure 4.2.6 | Distribution of ENC in different species in pisces, aves and mammals for MT-COIII gene | 98 |
| Figure 4.2.7 | Distribution of CAI in different species of pisces, aves and mammals for COI gene | 99 |
| Figure 4.2.8 | Distribution of CAI in different species of pisces, aves and mammals for COII gene | 100 |
| Figure 4.2.9 | Distribution of CAI in different species of pisces, aves and mammals for COIII gene | 100 |
| Figure 4.2.10 | Nucleotide composition and its composition at 3 rd codon position for MT-COI gene | 101 |
| Figure 4.2.11 | GC content and its content at codon's 1 st , 2 nd and 3 rd position in pisces, aves and mammals for MT-COI gene | 102 |
| Figure 4.2.12 | Nucleotide composition and its composition at 3 rd codon position for COII gene | 103 |
| Figure 4.2.13. | GC content and its content at codon's 1 st , 2 nd and 3 rd position in pisces, aves and mammals for MT-COII gene | 104 |
| Figure 4.2. 14. | Nucleotide composition and its composition at 3 rd codon position for COIII gene | 104 |

| | | |
|-------------------|---|-----|
| Figure 4.2.15 | GC content and its content at codon's 1st, 2 nd and 3 rd position in pisces, aves and mammals for MT-COIII gene | 105 |
| Figure 4.2.16 | Neutrality plot of GC12 with GC3 in different species of pisces for COI gene | 106 |
| Figure 4.2.16 (b) | Neutrality plot of GC12 with GC3 in different species of aves for COI gene | 108 |
| Figure 4.2.16 (c) | Neutrality plot of GC12 with GC3 in different species of mammals for COI gene | 108 |
| Figure 4.2.17(a) | Neutrality plot of GC12 with GC3 in different species of pisces for COII gene | 109 |
| Figure 4.2.17(b) | Neutrality plot of GC12 with GC3 in different species of aves for COII gene | 110 |
| Figure 4.2.17(c) | Neutrality plot of GC12 with GC3 in different species of mammals for COII gene | 111 |
| Figure 4.2.18 (a) | Neutrality plot of GC12 with GC3 in different species of pisces for COIII gene | 111 |
| Figure 4.2.18 (b) | Neutrality plot of GC12 with GC3 in different species of aves for COIII gene | 113 |
| Figure 4.2.18(c) | Neutrality plot of GC12 with GC3 in different species of mammals for COIII gene | 113 |
| Figure 4.2.19 | Contributions of the axes for MT-COI gene in pisces, aves and mammals | 113 |
| Figure 4.2.20 | Correspondence analysis of the synonymous codon usage for COI gene in pisces, aves and mammals | 114 |
| Figure 4.2.21 | Contributions of the axes for MT-COII gene in pisces, aves and mammals | 114 |
| Figure 4.2.22 | Correspondence analysis of the synonymous codon usage for COII gene in pisces, aves and mammals | 115 |
| Figure 4.2.23 | Contributions of the axes for MT-COIII gene in pisces, aves and mammals | 115 |
| Figure 4.2.25 | Correspondence analysis of the synonymous codon usage for COIII gene in pisces, aves and mammals | 116 |

| | | |
|-----------------|--|-----|
| Figure 4.2.26 | Comparison of amino acids in different species of pisces, aves and mammals for COI gene | 116 |
| Figure 4.2.27 | Comparison of amino acids in different species of pisces, aves and mammals for COII gene | 117 |
| Figure 4.2.28 | Comparison of amino acids in different species of pisces, aves and mammals for COIII gene | 117 |
| Figure 4.3.1 | Hierarchical Clustering of RSCU values of each codon among pisces, aves and mammals for CYB gene | 118 |
| Figure 4.3.2 | Distribution of ENC in different species of pisces, aves and mammals for MT-CYB gene | 118 |
| Figure 4.3.3 | Expression level in different species of pisces, aves and Mammals for MT-CYB gene | 119 |
| Figure 4.3.4 | Nucleotide composition and its 3 rd codon position for MT-CYB gene | 120 |
| Figure 4.3.5 | GC content and its content at codon's 1 st , 2 nd and 3 rd position in pisces, aves and mammals for MT-CYB gene | 121 |
| Figure 4.3.6(a) | Neutrality plot of GC12 with GC3 in different species of pisces for CYB gene | 121 |
| Figure 4.3.6(b) | Neutrality plot of GC12 with GC3 in different species of aves for CYB gene | 124 |
| Figure 4.3.6(c) | Neutrality plot of GC12 with GC3 in different species of mammals for CYB gene | 125 |
| Figure 4.3.7 | Contributions of the axes for MT-CYB in pisces, aves and mammals | 125 |
| Figure 4.3.8 | Correspondence analysis of the synonymous codon usage for CYB gene in pisces, aves and mammals | 125 |
| Figure 4.3.9 | Comparison of amino acids in different species of pisces, aves and mammals in CYB protein | 126 |
| Figure 4.4.1 | Hierarchical Clustering of RSCU values of each codon among pisces, aves and mammals for ATP6 | 127 |
| Figure 4.4.2 | Hierarchical Clustering of RSCU values of each codon among pisces, aves and mammals for MT-ATP8 | 127 |
| Figure 4.4.3 | Distribution of ENC in different species of pisces, aves and mammals for MT-ATP6 | 128 |

| | | |
|------------------|--|-----|
| Figure 4.4.4 | ENC distribution in different species of pisces, aves and mammals for ATP 8 | 129 |
| Figure 4.4.5 | Expression level in different species of pisces, aves and Mammals for MT-ATP6 | 129 |
| Figure 4.4.6 | Expression level in different species of pisces, aves and mammals for MT-ATP8 | 130 |
| Figure 4.4.7 | Nucleotide composition and its 3 rd codon position for MT-ATP6 | 130 |
| Figure 4.4.8 | Overall GC content and its content at codon's 1 st , 2 nd and 3 rd position in pisces, aves and mammals | 131 |
| Figure 4.4.9 | Nucleotide composition and its composition at 3 rd codon position for ATP8 gene | 132 |
| Figure 4.4.10 | Overall GC% and its content at 1 st , 2 nd and 3 rd codon position for ATP8 gene | 132 |
| Figure 4.4.11(a) | Neutrality plot of GC12 with GC3 in different species of pisces for ATP 6 gene | 133 |
| Figure 4.4.11(b) | Neutrality Plot of GC12 with GC3 in different species of aves for ATP 6 gene | 135 |
| Figure 4.4.11(c) | Neutrality plot of GC12 with GC3 in different species of mammals for ATP gene | 136 |
| Figure 4.4.12(a) | Neutrality plot of GC12 with GC3 in different species of pisces for ATP8 gene | 137 |
| Figure 4.4.12(b) | Neutrality plot of GC12 with GC3 in different species of aves for ATP8 gene | 138 |
| Figure 4.4.12(c) | Neutrality plot of GC12 with GC3 in different species of mammals for ATP8 gene | 138 |
| Figure 4.4.13 | Contributions of the axes for MT-ATP6 in pisces, aves and mammals | 139 |
| Figure 4.4.14 | Correspondence analysis of the synonymous codon usage for ATP6 gene in pisces, aves and mammals | 139 |
| Figure 4.4.15 | Contributions of the axes for MT-ATP8 gene in pisces, aves and mammals | 140 |

| | | | |
|--|---------------|---|-----|
| | Figure 4.4.16 | Correspondence analysis of the synonymous codon usage for ATP8 gene in pisces, aves and mammals | 140 |
| | Figure 4.4.17 | Comparison of amino acids in different species of pisces, aves and mammals for ATP6 gene | 141 |
| | Figure 4.4.18 | Comparison of amino acids in different species of pisces, aves and mammals for ATP8 gene | 141 |